

Original Article

Cite this article: Strauss P, Cook A, Winter S, Watson V, Wright Toussaint D, Lin A (2020). Associations between negative life experiences and the mental health of trans and gender diverse young people in Australia: findings from Trans Pathways. *Psychological Medicine* 50, 808–817. <https://doi.org/10.1017/S0033291719000643>

Received: 3 August 2018

Revised: 26 February 2019

Accepted: 7 March 2019

First published online: 8 July 2019

Key words:

Gender diversity; mental health; suicide; TGD; transgender

Author for correspondence:

Penelope Strauss, E-mail: penelope.strauss@telethonkids.org.au

Associations between negative life experiences and the mental health of trans and gender diverse young people in Australia: findings from Trans Pathways

Penelope Strauss¹, Angus Cook², Sam Winter³, Vanessa Watson⁴, Dani Wright Toussaint⁵ and Ashleigh Lin⁵

¹Telethon Kids Institute & School of Population and Global Health, University of Western Australia, Perth, Australia;

²School of Population and Global Health, University of Western Australia, Perth, Australia; ³School of Public Health, Curtin University, Perth, Australia; ⁴Western Australian Department of Health, YouthLink, North Metropolitan Area

Health Service, Perth, Australia and ⁵Telethon Kids Institute, University of Western Australia, Perth, Australia

Abstract

Background. Trans and gender diverse (TGD) young people worldwide experience high rates of poor mental health; however, these rates were unknown in Australia. In addition, how negative life events affect the mental health of TGD young people has been largely unexplored.

Methods. This paper reports on novel mental health findings of Trans Pathways, the largest study ever conducted in Australia with trans (transgender) and gender diverse young people ($N=859$; aged 14–25 years). The study was an anonymous online cross-sectional survey undertaken in 2016. Logistic and linear regression models were used to test associations between mental health outcomes and negative life experiences.

Results. TGD young people in Australia experience high levels of mental distress, including self-harming (79.7%), suicidal thoughts (82.4%), and attempting suicide (48.1%). Three in four participants had been diagnosed with depression and/or anxiety (74.6% and 72.2%, respectively). Many TGD young people had been exposed to negative experiences such as peer rejection (89.0%), precarious accommodation (22.0%), bullying (74.0%), and discrimination (68.9%). Most poor mental health outcomes were associated with negative experiences. The strongest associations were found for precarious accommodation and issues within educational settings. For example, participants with a prior suicide attempt were almost six times more likely to have experienced issues with accommodation, including homelessness.

Conclusions. The current results highlight the urgent need for better mental health care and provide insight into areas for targeted mental health interventions. These findings are pertinent for clinicians working with trans young people and wider society.

Introduction

People who are trans (transgender) or gender diverse have a gender incongruent with the sex assigned to them at birth. In recent years, there has been an increase in the prevalence of trans and gender diverse (TGD) young people, evidenced through population-based studies and the higher number of young people seeking gender-affirming interventions at gender clinics worldwide (Cohen-Kettenis *et al.*, 2011; Telfer *et al.*, 2015). The number of trans and gender non-conforming young people in the population is currently estimated to be almost 2.7% of adolescents (Rider *et al.*, 2018), higher than previous population estimates of 0.5–1.2% (Telfer *et al.*, 2015; Winter *et al.*, 2016; Flores *et al.*, 2017).

Research on the mental health of TGD populations tends to be conflated with research into LGBTIQ (lesbian, gay, bisexual, trans, intersex and queer) groups broadly. However, gender and sexuality are two very distinct aspects of a person's identity. LGBT (lesbian, gay, bisexual and trans) young people do experience poorer mental health than the general population, but these rates of poor mental health are even higher rates for trans youth compared to their lesbian, gay and bisexual peers (Hillier *et al.*, 2010; Veale *et al.*, 2017). It has been established that trans populations experience higher rates of poor mental health than their cisgender peers – those whose gender identity matches the sex assigned to them at birth (Bouman *et al.*, 2017; Veale *et al.*, 2017; Rider *et al.*, 2018). Previous research shows that children who are gender non-conforming in childhood are more likely to experience depression through adolescence and young adulthood compared to those who are gender conforming, in part attributable to adverse life events such as increased exposure to bullying and child abuse (Roberts *et al.*, 2013). Reisner *et al.* (2015) found that, in comparison to cisgender controls, trans young people experienced a two to threefold greater risk of anxiety disorders and depression.

Suicide attempts are relatively common in young trans populations – 37.8% of trans young people (aged 19–25) in Canada and 45% of trans adolescents in the UK have attempted suicide (Bradlow *et al.*, 2017; Veale *et al.*, 2017). TGD young people are significantly more likely than cisgender youth (regardless of sexuality) to self-harm and attempt suicide (Jones and Hillier, 2013; Smith *et al.*, 2014; Reisner *et al.*, 2015; Testa *et al.*, 2017). For example, trans young people are more likely to experience suicidal thoughts (31% compared to 11%) and to attempt suicide (17.2% compared to 6.1%) in comparison to matched cisgender controls (Reisner *et al.*, 2015). Causes of suicidal ideation and risk in TGD populations are likely to be multifactorial, including being attributable to minority stress, transphobia and life crises (McDermott *et al.*, 2017; Testa *et al.*, 2017). A previous national study of the mental health of TGD young people in Australia ($n = 189$) reported that, of the participants who had been exposed to abuse or discrimination, 70% had self-harmed and 37% had attempted suicide (Smith *et al.*, 2014). A history of physical and/or sexual violence has been associated with suicidal thoughts and attempts in adult trans populations (Testa *et al.*, 2012).

There are multiple and interacting factors contributing to mental health difficulties in this population. Previous research has indicated that trans populations may experience additional stress in a variety of situations (such as bathrooms and identification documents not matching gender expression) stemming from the expectation of rejection (Rood *et al.*, 2016). LGBT youth who experience peer victimisation experience higher levels of depressive symptoms, and have a lower sense of belonging within schools (Collier *et al.*, 2013). As a young person's gender non-conformity increases, so too do their likelihood of being bullied (Gordon *et al.*, 2018).

There has been limited research on the mental health of TGD young people within Australia. In particular, the impact of negative and/or traumatic events on mental wellbeing in this population has not been extensively examined (Reisner *et al.*, 2016). The only previous national study of Australian TGD young people reported on some aspects of mental health but did not investigate the associations between potential drivers of poor mental health, such as discrimination and bullying, and adverse health outcomes; e.g. self-harm, suicidality and psychiatric diagnoses (Smith *et al.*, 2014). The earlier study focussed on suicide and self-harm after exposure to violence, but not wider patterns and predictors. This limits our understanding of why TGD young people experience elevated rates of mental health difficulties and constrains our ability to develop and implement appropriate intervention strategies (Reisner *et al.*, 2016). This study aims to characterise mental health issues affecting TGD Australian young people, and to investigate the potential relationships between negative life events and adverse mental health outcomes.

Methods

Study population

The survey sample comprised 859 young people aged 14–25 years who self-identified as trans or gender diverse (TGD) and were currently residing in Australia between February and August 2016.

Study design

An online cross-sectional survey was undertaken. TGD young people and parents of TGD young people were consulted to

determine their preferences for questions to be included in the survey, and to ensure that the questions were relevant and asked in a respectful manner. We held a focus group with TGD young people and a separate focus group with parents of TGD young people. These groups were provided with an initial draft of the questionnaire and the final version was shaped by their feedback from these sessions. The focus group members identified drivers and protective factors of mental health based on their own experiences and awareness of concerns raised within the community. Qualtrics online survey software was used to construct and host the questionnaire which utilised branch, display and skip logic based on participant responses. All questions were optional, except those used to determine eligibility (i.e. TGD identification, age, place of residence).

Recruitment and consent procedures

An anonymous online, self-report questionnaire was conducted between February and August 2016. Participants were largely recruited using social media (i.e. Twitter, Facebook and Tumblr), gender clinics, youth mental health services, support groups, parent and youth groups, and word of mouth. Participants were provided with an online participant information sheet and were instructed that by entering the online survey they were consenting to take part in the study. Parental consent was not required. The study was approved by the University of Western Australia ethics committee (RA/4/1/7958).

Outcomes and risk factors of interest

The questionnaire included both quantitative and qualitative components. Primary outcomes of interest were self-reported psychiatric diagnoses, adverse health outcomes, and current anxiety and depressive symptomatology. Exposures to a range of negative life events and stressors were also assessed.

Gender and demographics

Participants were asked for both their sex assigned at birth (male/female) and gender identity (open text box). Asking about gender in this way allowed participants to describe their gender in their own words. The study was advertised as a study for TGD young people. Participants were asked for their year of birth and whether they were living in Australia at the time of the survey to determine their eligibility. Participants were also asked about their current living situation.

Current psychopathology

Depressive symptoms were indexed on the Patient Health Questionnaire (PHQ-A) (Kroenke *et al.*, 2001). The PHQ-A is a nine-item scale that is scored from 0–27 and can be categorised into no depressive disorder (0–4); possible mild depressive disorder (5–9); possible moderate depressive disorder (10–14); possible moderately severe depressive disorder (15–19); and possible severe depressive disorder (20–27) (Kroenke *et al.*, 2001). Anxiety was measured using the Generalised Anxiety Disorder seven-item Scale (GAD-7) (Spitzer *et al.*, 2006). The categories for the GAD-7 are based on scores suggestive of a generalised anxiety disorder (≥ 5); moderate to severe anxiety (≥ 10); and severe anxiety (≥ 15) (Spitzer *et al.*, 2006).

Table 1. Lifetime prevalence of self-harm, suicidal thoughts and suicide attempts among trans young people

Self-reported adverse health outcome	Group identification				
	Overall sample: <i>n</i> (%)	Assigned female at birth: <i>n</i> (%)	Assigned male at birth: <i>n</i> (%)	Aged <18: <i>n</i> (%)	Aged ≥18: <i>n</i> (%)
Desire to self-harm	639 (91.3)	493 (93.5)**	146 (84.4)**	174 (91.1)	465 (91.4)
Self-harming	561 (79.7)	446 (84.6)**	115 (65.0)**	149 (78.0)	412 (80.3)
Reckless behaviour to purposely put life at risk	432 (62.8)	336 (65.5)*	96 (54.9)*	104 (59.8)	328 (63.8)
Suicidal thoughts	568 (82.4)	435 (84.1)*	133 (77.3)*	147 (83.1)	421 (82.2)
Suicide attempt	333 (48.1)	258 (49.8)	75 (43.1)	77 (42.1)	256 (50.3)

* χ^2 test significant difference between the two comparative groups ($p < 0.05$).

** χ^2 test significant difference between the two comparative groups ($p < 0.01$).

Self-reported psychiatric diagnoses

Psychiatric diagnoses (depression, anxiety disorders, post-traumatic stress disorder, eating disorders, autism spectrum disorder, personality disorders, psychosis and substance use disorders) were listed and participants were asked whether a health professional had ever diagnosed the individual with the specific psychiatric conditions.

Self-reported adverse health outcomes

The five self-reported adverse health outcomes measured were: wanting to self-harm, self-harming, reckless behaviour that purposely puts one's life at risk, suicidal thoughts and suicide attempts. These were measured by asking participants whether they had engaged in the outcome within the last 12 months, prior to the last 12 months or never. Here we report a lifetime prevalence of these adverse health outcomes.

Exposure to negative experiences

Participants were asked about a range of negative experiences that are potentially associated with poor mental health. Participants were asked to select all the factors that they had experienced from the list provided. These items included: issues with accommodation (including homelessness), body dysphoria, bullying, discrimination, employment issues, experiencing a significant loss, feeling isolated from not knowing other TGD people, feeling isolated from services, helping others with their issues with mental health, a lack of family support, peer rejection and issues with school, university or technical college (henceforth education settings).

Statistical analysis

IBM SPSS Statistics, version 24, and Stata, version 15, were used to obtain descriptive statistics (frequencies, means, standard deviations), and to develop regression models. Logistic regression models were used to evaluate associations between potential drivers and adverse health outcomes, psychiatric diagnoses, and current psychopathology using known cut off-scores on the PHQ-A and the GAD-7. Linear regression models were used to evaluate potential drivers of poor mental health and current psychopathology measured by the GAD-7 and the PHQ-A. All reported regressions are adjusted for age and sex assigned at birth. All

regression models were evaluated using diagnostic testing and no major deviations in distributional assumptions were detected.

Results

Demographics

The majority of participants were assigned female at birth (74.4%). Aboriginal and/or Torres Strait Islander people comprised 3.7% of this sample, a proportion that is representative of Australian population demographics (Australian Bureau of Statistics, 2014). The mean age of participants was 19.37 years (*s.d.* = 3.15). A total of 29.7% ($n = 255$) of participants self-identified as trans male or male, 15% ($n = 129$) identified as trans female or female, and 48.6% ($n = 417$) identified as various non-binary identities including non-binary transmasculine, non-binary femme, agender, bigender, pangender and other non-binary identities.

Most participants were living with parents (60.3%), 18.7% were in shared accommodation, 4.7% were living with other family members, 4.3% were living with a partner, 3.7% were living alone, 3.4% were in a residential college, 2.3% in supported accommodation and 1.4% had no fixed accommodation. One quarter of participants lived in Victoria (25.2%), 20.0% in New South Wales, 17.2% in Queensland, 15.9% in Western Australia, 12.0% in South Australia, 6.5% in the Australian Capital Territory, 2.7% in Tasmania and 0.5% in the Northern Territory. There was slight overrepresentation in the Australian Capital Territory, South Australia, Tasmania, Victoria and Western Australia while New South Wales, the Northern Territory, Queensland were underrepresented compared to 2015 population estimates by the Australian Bureau of Statistics (Australian Bureau of Statistics, 2016).

Self-harm and suicidality

Table 1 reports the lifetime prevalence rates of adverse health outcomes among our sample. Self-harm was commonly reported, with 91.3% ever wanting to self-harm and 79.7% self-harming at some point during their life. A majority of participants had engaged in reckless behaviour to risk their life (62.8%). Over three quarters (82.4%) of participants reported ever having suicidal thoughts and 48.1% had ever attempted suicide. There were no statistically significant differences in the prevalence of all lifetime adverse health outcomes between TGD young people




Fig. 1. Prevalence of self-reported psychiatric diagnoses among trans young people ($n = 756$).

under the age of 18 compared to those aged 18 or older. However, there were significant differences between TGD young people who were birth-assigned female compared to those birth-assigned male as tested with a Pearson χ^2 test. Participants assigned female at birth reported higher rates of wanting to self-harm ($F = 13.724$, $p < 0.001$), self-harming ($F = 31.633$, $p < 0.001$), reckless behaviour to purposely put one's life at risk ($F = 6.323$, $p = 0.012$) and suicidal thoughts ($F = 4.139$, $p = 0.042$). There was no statistically significant difference between these groups for suicide attempts.

Self-reported psychiatric diagnoses

Figure 1 summarises the prevalence of self-reported psychiatric diagnoses ever received by a health professional. The most prevalent diagnoses were depression (74.6%) followed by anxiety (72.2%) and post-traumatic stress disorder (25.1%). There were no statistically significant differences in prevalence between participants assigned female at birth and those assigned male at birth across all diagnoses.

Current psychopathology

More participants scored in the moderate to severe anxiety range with 11.8% of participants reporting minimal anxiety, 26.0% reporting mild anxiety, 30.5% reporting moderate anxiety and 31.6% reporting severe anxiety during the prior 2 weeks ($n = 845$). The mean GAD-7 score was 11.53 (s.d. = 5.65, $n = 845$). Based on the GAD-7 functional impairment scale, only 7.6% of the sample ($n = 65$) had no functional impairment, while 52.2% ($n = 443$) had some difficulties, 28.0% ($n = 238$) found it very difficult to carry out everyday tasks and 12.3% ($n = 105$) reported extreme difficulties. Depressive symptoms similarly increased in severity with 7.5% of participants reporting no depressive symptoms, 16.2% with mild depressive symptoms, 21.6% with moderate depressive symptoms, 24.6% with moderately severe depressive

symptoms and 30.2% with severe depressive symptoms during the 2 weeks prior to completing the survey ($n = 736$). The mean PHQ-A score was 15.26 (s.d. = 7.04, $n = 736$).

Exposure to negative experiences

Young people in this population were commonly exposed to negative experiences, including peer rejection (89.0%), issues within educational settings (78.9%) and bullying (74.0%). Body dysphoria was commonly experienced (93.8%). These patterns of exposure are summarised in Tables 2–4.

Associations between negative experiences and self-harm and suicidality

Table 2 reports the associations between potential drivers of poor mental health and the five adverse health outcomes of self-harm and suicide measured. The majority were associated with an adverse outcome. For example, participants with a history of self-harm had significantly elevated odds ratios for exposure to issues with accommodation (including homelessness) (OR 4.099, 95% CI 2.131–7.883) and within educational settings (OR 3.539, 95% CI 2.301–5.442). Participants with a prior suicide attempt had significantly elevated odds ratios for exposure to issues with accommodation (OR 5.716, 95% CI 3.617–9.031) and within educational settings (OR 3.892, 95% CI 2.528–5.992).

Associations between negative experiences and psychiatric diagnoses

Table 3 summarises the associations between potential drivers of poor mental health and psychiatric diagnoses. Participants who had been diagnosed with depression reported a greater than threefold increase in their likelihood of experiencing issues within educational settings (OR 3.604, 95% CI 2.424–5.359) and

Table 2. Potential predictors of self-harming and suicidal behaviours

Personal event/ experiences reported	Overall sample reported: <i>n</i> (%)	Lifetime desire to self-harm			Lifetime self-harming			Lifetime engaging in reckless behaviour (to purposely put life at risk)			Lifetime suicidal thoughts			Lifetime suicide attempt		
		Outcome Y/N (total)	Adjusted odds ratio	<i>p</i> value	Outcome Y/N (total)	Adjusted odds ratio	<i>p</i> value	Outcome Y/N (total)	Adjusted odds ratio	<i>p</i> value	Outcome Y/N (total)	Adjusted odds ratio	<i>p</i> value	Outcome Y/N (total)	Adjusted odds ratio	<i>p</i> value
Accommodation issues	22.0%, <i>n</i> = 147	137/4 (<i>n</i> = 634)	5.101 (1.780–14.618)	<i>p</i> = 0.002**	132/12 (<i>n</i> = 637)	4.099 (2.131–7.883)	<i>p</i> < 0.001**	116/25 (<i>n</i> = 622)	4.103 (2.514–6.695)	<i>p</i> < 0.001**	139/4 (<i>n</i> = 625)	11.279 (4.030–31.563)	<i>p</i> < 0.001**	108/32 (<i>n</i> = 626)	5.716 (3.617–9.031)	<i>p</i> < 0.001**
Body dysphoria	93.8%, <i>n</i> = 648	561/55 (<i>n</i> = 656)	0.751 (0.222–2.542)	<i>p</i> = 0.645	497/121 (<i>n</i> = 659)	1.784 (0.878–3.625)	<i>p</i> = 0.109	385/217 (<i>n</i> = 642)	3.262 (1.657–6.432)	<i>p</i> = 0.001**	507/11 (<i>n</i> = 645)	2.634 (1.297–5.350)	<i>p</i> = 0.007**	296/309 (<i>n</i> = 646)	1.842 (0.941–3.605)	<i>p</i> = 0.074
Bullying	74.0%, <i>n</i> = 497	450/25 (<i>n</i> = 639)	5.048 (2.805–9.085)	<i>p</i> < 0.001**	404/74 (<i>n</i> = 640)	3.036 (1.978–4.662)	<i>p</i> < 0.001**	320/146 (<i>n</i> = 626)	2.932 (2.015–4.266)	<i>p</i> < 0.001**	413/53 (<i>n</i> = 630)	4.081 (2.628–6.338)	<i>p</i> < 0.001**	262/208 (<i>n</i> = 631)	3.664 (2.447–5.486)	<i>p</i> < 0.001**
Discrimination	68.9%, <i>n</i> = 454	402/25 (<i>n</i> = 625)	2.778 (1.573–4.906)	<i>p</i> < 0.001**	367/69 (<i>n</i> = 628)	2.111 (1.393–3.200)	<i>p</i> < 0.001**	301/119 (<i>n</i> = 614)	3.288 (2.298–4.703)	<i>p</i> < 0.001**	374/51 (<i>n</i> = 615)	2.847 (1.846–4.392)	<i>p</i> < 0.001**	238/187 (<i>n</i> = 617)	2.735 (1.905–3.927)	<i>p</i> < 0.001**
Employment issues	41.9%, <i>n</i> = 281	249/22 (<i>n</i> = 639)	1.571 (0.848–2.912)	<i>p</i> = 0.151	229/40 (<i>n</i> = 642)	2.235 (1.394–3.583)	<i>p</i> = 0.001**	198/70 (<i>n</i> = 625)	2.856 (1.943–4.199)	<i>p</i> < 0.001**	235/30 (<i>n</i> = 627)	2.515 (1.528–4.139)	<i>p</i> < 0.001**	163/105 (<i>n</i> = 635)	2.849 (1.972–4.118)	<i>p</i> < 0.001**
Experiencing a significant loss	53.3%, <i>n</i> = 359	328/14 (<i>n</i> = 639)	3.835 (2.023–7.271)	<i>p</i> < 0.001**	290/51 (<i>n</i> = 644)	1.836 (1.222–2.757)	<i>p</i> = 0.003**	240/99 (<i>n</i> = 627)	2.154 (1.543–3.007)	<i>p</i> < 0.001**	299/42 (<i>n</i> = 630)	2.119 (1.381–3.250)	<i>p</i> = 0.001**	185/147 (<i>n</i> = 633)	1.830 (1.327–2.524)	<i>p</i> < 0.001**
Feeling isolated from not knowing other trans people	66.1%, <i>n</i> = 455	403/31 (<i>n</i> = 652)	1.700 (0.975–2.965)	<i>p</i> = 0.062	354/79 (<i>n</i> = 654)	1.331 (0.884–2.004)	<i>p</i> = 0.171	278/146 (<i>n</i> = 639)	1.439 (1.022–2.026)	<i>p</i> = 0.037*	357/69 (<i>n</i> = 641)	1.271 (0.828–1.950)	<i>p</i> = 0.272	210/210 (<i>n</i> = 641)	1.275 (0.915–1.775)	<i>p</i> = 0.151
Feeling isolated from services	60.1%, <i>n</i> = 404	360/28 (<i>n</i> = 640)	1.684 (0.971–2.919)	<i>p</i> = 0.064	327/62 (<i>n</i> = 644)	1.870 (1.253–2.792)	<i>p</i> = 0.002**	255/121 (<i>n</i> = 625)	1.887 (1.354–2.631)	<i>p</i> < 0.001**	329/54 (<i>n</i> = 629)	1.804 (1.191–2.734)	<i>p</i> = 0.005**	197/181 (<i>n</i> = 628)	1.586 (1.146–2.196)	<i>p</i> = 0.005**
Helping others with their issues with mental health	70.2%, <i>n</i> = 473	430/23 (<i>n</i> = 642)	3.249 (1.818–5.806)	<i>p</i> < 0.001**	387/70 (<i>n</i> = 645)	2.264 (1.495–3.428)	<i>p</i> < 0.001**	285/152 (<i>n</i> = 628)	1.465 (1.029–2.086)	<i>p</i> = 0.034*	381/61 (<i>n</i> = 630)	2.140 (1.391–3.293)	<i>p</i> = 0.001**	217/224 (<i>n</i> = 633)	1.130 (0.800–1.595)	<i>p</i> = 0.488
Lack of family support	65.8%, <i>n</i> = 431	382/26 (<i>n</i> = 620)	2.353 (1.340–4.132)	<i>p</i> = 0.003**	344/70 (<i>n</i> = 624)	1.899 (1.263–2.856)	<i>p</i> = 0.002**	269/132 (<i>n</i> = 610)	1.979 (1.401–2.795)	<i>p</i> < 0.001**	354/52 (<i>n</i> = 610)	2.547 (1.662–3.902)	<i>p</i> < 0.001**	216/190 (<i>n</i> = 616)	1.913 (1.358–2.696)	<i>p</i> < 0.001**
Peer rejection	89.0%, <i>n</i> = 613	534/44 (<i>n</i> = 651)	2.731 (1.378–5.413)	<i>p</i> = 0.004**	478/108 (<i>n</i> = 656)	2.602 (1.502–4.507)	<i>p</i> = 0.001**	367/202 (<i>n</i> = 638)	2.282 (1.371–3.798)	<i>p</i> = 0.001**	482/88 (<i>n</i> = 639)	2.610 (1.494–4.560)	<i>p</i> = 0.001**	285/289 (<i>n</i> = 643)	2.124 (1.242–3.630)	<i>p</i> = 0.006**
School, university or TAFE issues	78.9%, <i>n</i> = 542	486/28 (<i>n</i> = 651)	4.505 (2.550–7.959)	<i>p</i> < 0.001**	437/78 (<i>n</i> = 654)	3.539 (2.301–5.442)	<i>p</i> < 0.001**	345/156 (<i>n</i> = 637)	3.592 (2.414–5.347)	<i>p</i> < 0.001**	439/63 (<i>n</i> = 640)	3.488 (2.240–5.431)	<i>p</i> < 0.001**	275/230 (<i>n</i> = 644)	3.892 (2.528–5.992)	<i>p</i> < 0.001**

Odds ratios adjusted for sex assigned at birth and age (by year of birth).

p* < 0.05; *p* < 0.01.

Table 3. Psychiatric diagnoses and associations with negative experiences

Personal event/ experiences reported	Overall sample reported: <i>n</i> (%)	Lifetime diagnosis of depression	Lifetime diagnosis of anxiety	Lifetime diagnosis of PTSD	Lifetime diagnosis of a personality disorder	Lifetime diagnosis of psychosis	Lifetime diagnosis of an eating disorder	Lifetime diagnosis of ASD	Lifetime diagnosis of a substance use disorder
Accommodation issues	22.0%, <i>n</i> = 147	3.214 (1.796–5.752) <i>p</i> < 0.001**	2.653 (1.586–4.439) <i>p</i> < 0.001**	3.285 (2.173–4.965) <i>p</i> < 0.001**	2.309 (1.481–3.600) <i>p</i> < 0.001**	2.641 (1.655–4.214) <i>p</i> < 0.001**	2.487 (1.612–3.837) <i>p</i> < 0.001**	1.883 (1.216–2.916) <i>p</i> = 0.005**	1.755 (1.055–2.921) <i>p</i> = 0.030**
Body dysphoria	93.8%, <i>n</i> = 648	1.465 (0.745–2.882) <i>p</i> = 0.268	1.109 (0.562–2.188) <i>p</i> = 0.766	1.482 (0.668–3.285) <i>p</i> = 0.333	1.946 (0.750–5.044) <i>p</i> = 0.171	2.002 (0.700–5.724) <i>p</i> = 0.195	1.915 (0.791–4.633) <i>p</i> = 0.150	0.841 (0.413–1.714) <i>p</i> = 0.633	1.287 (0.492–3.362) <i>p</i> = 0.607
Bullying	74.0%, <i>n</i> = 497	2.678 (1.827–3.924) <i>p</i> < 0.001**	1.880 (1.292–2.737) <i>p</i> = 0.001**	1.904 (1.215–2.984) <i>p</i> = 0.005**	2.106 (1.279–3.468) <i>p</i> = 0.003**	1.834 (1.077–3.123) <i>p</i> = 0.026*	2.154 (1.348–3.442) <i>p</i> = 0.001**	1.236 (0.800–1.908) <i>p</i> = 0.340	1.166 (0.688–1.975) <i>p</i> = 0.569
Discrimination	68.9%, <i>n</i> = 454	1.476 (1.013–2.151) <i>p</i> = 0.043*	1.485 (1.031–2.138) <i>p</i> = 0.034*	2.003 (1.318–3.045) <i>p</i> = 0.001**	2.298 (1.436–3.679) <i>p</i> = 0.001**	1.757 (1.080–2.858) <i>p</i> = 0.023*	2.436 (1.554–3.817) <i>p</i> < 0.001**	1.405 (0.926–2.132) <i>p</i> = 0.110	1.744 (1.031–2.952) <i>p</i> = 0.038*
Employment issues	41.9%, <i>n</i> = 281	1.923 (1.265–2.925) <i>p</i> = 0.002**	1.759 (1.181–2.621) <i>p</i> = 0.005**	1.401 (0.941–2.086) <i>p</i> = 0.097	1.621 (1.054–2.492) <i>p</i> = 0.028*	1.622 (1.017–2.587) <i>p</i> = 0.042*	1.767 (1.168–2.671) <i>p</i> = 0.007**	1.068 (0.702–1.623) <i>p</i> = 0.759	1.435 (0.875–2.352) <i>p</i> = 0.153
Experiencing a significant loss	53.3%, <i>n</i> = 359	2.245 (1.555–3.242) <i>p</i> < 0.001**	1.569 (1.104–2.229) <i>p</i> = 0.012*	1.200 (0.840–1.714) <i>p</i> = 0.315	1.175 (0.802–1.722) <i>p</i> = 0.409	1.495 (0.981–2.276) <i>p</i> = 0.061	1.712 (1.177–2.491) <i>p</i> = 0.005**	1.203 (0.827–1.750) <i>p</i> = 0.333	1.286 (0.815–2.029) <i>p</i> = 0.280
Feeling isolated from not knowing other trans people	66.1%, <i>n</i> = 455	1.497 (1.040–2.155) <i>p</i> = 0.030*	1.192 (0.837–1.698) <i>p</i> = 0.330	1.597 (1.086–2.348) <i>p</i> = 0.017*	1.154 (0.774–1.721) <i>p</i> = 0.483	1.400 (0.897–2.185) <i>p</i> = 0.139	1.488 (1.007–2.200) <i>p</i> = 0.046*	1.564 (1.044–2.341) <i>p</i> = 0.030**	1.056 (0.667–1.672) <i>p</i> = 0.817
Feeling isolated from services	60.1%, <i>n</i> = 404	1.423 (0.993–2.037) <i>p</i> = 0.054	1.591 (1.125–2.250) <i>p</i> = 0.009**	1.767 (1.213–2.574) <i>p</i> = 0.003**	0.970 (0.658–1.432) <i>p</i> = 0.880	1.230 (0.803–1.884) <i>p</i> = 0.342	1.252 (0.863–1.817) <i>p</i> = 0.237	1.144 (0.784–1.667) <i>p</i> = 0.486	0.776 (0.499–1.207) <i>p</i> = 0.261
Helping others with their issues with mental health	70.2%, <i>n</i> = 473	1.387 (0.945–2.035) <i>p</i> = 0.095	1.261 (0.872–1.825) <i>p</i> = 0.218	1.044 (0.707–1.540) <i>p</i> = 0.830	1.187 (0.774–1.820) <i>p</i> = 0.432	0.942 (0.602–1.473) <i>p</i> = 0.792	1.020 (0.685–1.519) <i>p</i> = 0.923	0.896 (0.600–1.337) <i>p</i> = 0.590	0.740 (0.464–1.180) <i>p</i> = 0.206
Lack of family support	65.8%, <i>n</i> = 431	1.844 (1.275–2.666) <i>p</i> = 0.001**	1.900 (1.329–2.717) <i>p</i> < 0.001**	1.674 (1.126–2.489) <i>p</i> = 0.011*	1.529 (0.998–2.343) <i>p</i> = 0.051	1.202 (0.769–1.879) <i>p</i> = 0.419	1.956 (1.285–2.978) <i>p</i> = 0.002**	0.906 (0.615–1.333) <i>p</i> = 0.616	1.114 (0.688–1.804) <i>p</i> = 0.662
Peer rejection	89.0%, <i>n</i> = 613	1.629 (0.970–2.735) <i>p</i> = 0.065	1.641 (0.991–2.717) <i>p</i> = 0.054	1.089 (0.619–1.916) <i>p</i> = 0.767	1.248 (0.665–2.342) <i>p</i> = 0.489	1.042 (0.542–2.003) <i>p</i> = 0.901	1.561 (0.834–2.921) <i>p</i> = 0.163	1.194 (0.658–2.168) <i>p</i> = 0.560	0.796 (0.411–1.540) <i>p</i> = 0.796
School, university or TAFE issues	78.9%, <i>n</i> = 542	3.604 (2.424–5.359) <i>p</i> < 0.001**	3.162 (2.149–4.654) <i>p</i> < 0.001**	1.416 (0.900–2.228) <i>p</i> = 0.133	1.393 (0.852–2.278) <i>p</i> = 0.186	1.478 (0.860–2.541) <i>p</i> = 0.157	1.576 (0.981–2.532) <i>p</i> = 0.060	1.396 (0.873–2.233) <i>p</i> = 0.163	1.067 (0.621–1.834) <i>p</i> = 0.814

Odds ratios adjusted for sex assigned at birth and age (by year of birth).

p* < 0.05; *p* < 0.01.

Table 4. Self-reported current psychopathology and associations with negative experiences

Negative experiences reported	Overall sample reported: <i>n</i> (%)	Current severe anxiety (GAD-7 score = ≥ 15 v. < 15) OR (95% CI) <i>p</i> value	Current anxiety – linear regression β coefficient (95% CI) <i>p</i> value	Current depression (PHQ-A score = ≥ 15 v. < 15) OR (95% CI) <i>p</i> value	Current depression – linear regression β coefficient (95% CI) <i>p</i> value
Accommodation issues	22.0%, <i>n</i> = 147	2.379 (1.567–3.613) <i>p</i> < 0.001**	2.704 (1.677–3.731) <i>p</i> < 0.001**	3.374 (2.206–5.160) <i>p</i> < 0.001**	4.512 (3.253–5.771) <i>p</i> < 0.001**
Body dysphoria	93.8%, <i>n</i> = 648	1.206 (0.588–2.475) <i>p</i> = 0.610	1.417 (–0.304 to 3.139) <i>p</i> = 0.106	2.372 (1.210–4.649) <i>p</i> = 0.012*	1.880 (–0.278 to 4.037) <i>p</i> = 0.088
Bullying	74.0%, <i>n</i> = 497	1.988 (1.317–3.001) <i>p</i> = 0.001**	1.869 (0.912–2.823) <i>p</i> < 0.001**	2.454 (1.703–3.535) <i>p</i> < 0.001**	2.871 (1.692–4.050) <i>p</i> < 0.001**
Discrimination	68.9%, <i>n</i> = 454	1.566 (1.074–2.282) <i>p</i> = 0.020*	1.337 (0.424–2.250) <i>p</i> = 0.004**	1.896 (1.349–2.665) <i>p</i> < 0.001**	2.072 (0.956–3.188) <i>p</i> < 0.001**
Employment issues	41.9%, <i>n</i> = 281	1.785 (1.212–2.629) <i>p</i> = 0.003**	2.020 (1.086–2.954) <i>p</i> < 0.001**	2.353 (1.629–3.399) <i>p</i> < 0.001**	3.155 (2.000–4.310) <i>p</i> < 0.001**
Experiencing a significant loss	53.3%, <i>n</i> = 359	1.481 (1.054–2.082) <i>p</i> = 0.024*	1.376 (0.534–2.219) <i>p</i> = 0.001**	1.921 (1.395–2.647) <i>p</i> < 0.001**	2.545 (1.509–3.581) <i>p</i> < 0.001**
Feeling isolated from not knowing other trans people	66.1%, <i>n</i> = 455	1.526 (1.061–2.193) <i>p</i> = 0.023*	1.675 (0.810–2.539) <i>p</i> < 0.001**	2.161 (1.560–2.994) <i>p</i> < 0.001**	2.593 (1.530–3.656) <i>p</i> < 0.001**
Feeling isolated from services	60.1%, <i>n</i> = 404	1.684 (1.188–2.389) <i>p</i> = 0.003**	2.031 (1.190–2.872) <i>p</i> < 0.001**	2.428 (1.760–3.350) <i>p</i> < 0.001**	3.456 (2.426–4.486) <i>p</i> < 0.001**
Helping others with their issues with mental health	70.2%, <i>n</i> = 473	2.323 (1.549–3.484) <i>p</i> < 0.001**	2.319 (1.415–3.223) <i>p</i> < 0.001**	1.885 (1.339–2.654) <i>p</i> < 0.001**	2.732 (1.607–3.857) <i>p</i> < 0.001**
Lack of family support	65.8%, <i>n</i> = 431	1.683 (1.166–2.429) <i>p</i> = 0.005**	1.773 (0.879–2.667) <i>p</i> < 0.001**	1.692 (1.216–2.355) <i>p</i> = 0.002**	2.380 (1.281–3.480) <i>p</i> < 0.001**
Peer rejection	89.0%, <i>n</i> = 613	2.143 (1.162–3.954) <i>p</i> = 0.015*	2.606 (1.287–3.926) <i>p</i> < 0.001**	2.766 (1.655–4.622) <i>p</i> < 0.001**	3.629 (2.0141–5.243) <i>p</i> < 0.001**
School, university or TAFE issues	78.9%, <i>n</i> = 542	2.639 (1.633–4.266) <i>p</i> < 0.001**	2.779 (1.785–3.774) <i>p</i> < 0.001**	3.265 (2.197–4.851) <i>p</i> < 0.001**	4.618 (3.412–5.823) <i>p</i> < 0.001**

Odds ratios adjusted for sex assigned at birth and age (by year of birth). Current severe anxiety defined as scoring a 15 or above on the GAD-7 and current moderately severe to severe depressive symptoms defined as scoring 15 or above on the PHQ-A. **p* < 0.05; ***p* < 0.01.

accommodation issues (OR 3.214, 95% CI 1.796–5.752). For participants who had been diagnosed with an anxiety disorder, odds ratios of 3.2 (95% CI 2.149–4.654) were estimated for educational settings and 2.7 for accommodation issues (95% CI 1.586–4.439). Participants who had ever been diagnosed with PTSD were more than three times as likely to have experienced accommodation issues (OR 3.285, 95% CI 2.173–4.965) and twice as likely to have experienced discrimination (OR 2.003, 95% CI 1.318–3.045).

Associations between negative experiences and current psychopathology

Both issues with accommodation and issues within educational settings had the largest effect on all associations between life experiences and current psychopathology. Participants with severe current anxiety (scoring 15 or higher on the GAD-7) had more than a twofold increase of odds of exposure to accommodation issues (OR 2.379, 95% CI 1.567–3.613) and issues within educational settings (OR 2.639, 95% CI 1.633–4.266). The results were similar for participants with more severe current depressive symptoms (scoring 15 or higher on the PHQ-A) for exposure to accommodation issues (OR 3.374, 95% CI 2.206–5.160) and issues within educational settings (OR 3.265, 95% CI 2.197–4.851). These results are reflected in the linear regression modelling of the GAD-7 and PHQ-A instruments, as seen in Table 4.

Discussion

In the current study, we present the findings from a large sample of TGD young people and the indicators of their mental health. Results demonstrated high rates of current depressive and anxiety symptoms, and of self-reported psychiatric diagnoses. Rates of self-harm and suicidality were exceptionally high, with almost four in every five participants having a history of self-harm and nearly one in two having attempted suicide. Negative experiences, including experiencing unstable accommodation, discrimination, bullying, feeling unsupported from family, and issues in education and employment, were associated with most poor mental health outcomes in this sample. The largest associations between life experiences and self-harming and suicide attempts related to issues with housing and education. These are two areas that can be improved through interventions with families and educational environments to make them more supportive of TGD young people.

Self-harm and suicidality

The lifetime percentages of reported self-harm and suicide attempts were 79.7% and 48.1%, respectively. This suicide attempt rate is over 14 times greater than the general Australian adult population rate (Australian Bureau of Statistics, 2007). These rates are strikingly similar to those recently reported in the UK, where 84% of trans adolescents report self-harming and 45% report attempted suicide (Bradlow *et al.*, 2017). The lack of statistical difference between the rates of self-harm and suicidality between the under-18 and over-18 participants implies that self-harming and suicidal behaviours may tend to begin early and continue into young adulthood. Rates of suicidal ideation and suicide attempts are higher than rates seen internationally for trans adults, where a meta-synthesis averaged reported rates of suicidal ideation and attempts to be 55% and 29%, respectively (Adams *et al.*, 2017).

Mental health

Over three-quarters of participants reported exhibiting moderate to severe depressive symptoms during the previous 2 weeks (76.4%) and more than half of the participants reported moderate to severe anxiety during the previous 2 weeks (62.1%). These are consistent with the self-reported rates of diagnosed depression and anxiety. Current findings of depressive and anxiety symptomatology are higher than would be expected in the general population. Specifically, in comparison with the general Australian adolescent population, the rates of depressive symptoms are seven times higher and anxiety-related symptoms are over four times higher (Lawrence *et al.*, 2015). Our results similar to other literature that shows rates of depression, anxiety and emotional distress in trans young people to be higher than both the general population and same-gender attracted youth (Hillier *et al.*, 2010; Hyde *et al.*, 2014).

Co-occurring psychiatric disorders in gender diverse populations are unlikely to be simply related to the person's gender identity, and may instead reflect the response of the individual's familial and social environment to that identity and its expression (Vrouenraets *et al.*, 2015). It has been suggested that cisgender populations under the same incessant exposure to psychological stressors would in all likelihood experience comparable rates of depression and anxiety as trans populations (Inch, 2016). Higher rates of anxiety among transgender adult populations have also been reported, and have been attributed to low self-esteem and poor interpersonal functioning (Bouman *et al.*, 2017).

Exposure to negative experiences as potential drivers of poor mental health

Mental distress experienced by gender diverse populations does not intrinsically arise from the experience of an incongruent gender identity, but is more likely due to exposures to negative external events. Many of our participants were exposed to negative experiences known to be associated with poor mental health from other research (Mustanski and Liu, 2013; Roberts *et al.*, 2013; Taliaferro *et al.*, 2018), such as discrimination (68.9%), issues with accommodation (22.0%) and feeling isolated from other TGD people and services (66.1% and 60.1%, respectively). Notably, every potential driver of poor mental health was associated with participants reporting depressive symptoms at the more severe end of the spectrum. These risk factors are external (with the exception of body dysphoria) and are therefore potentially preventable. Previous research has indicated that prepubescent trans young people who are able to socially transition exhibit psychopathology similar to the general population. This suggests that if gender diverse children are supported to explore and affirm their gender identity they are more mentally healthy (Olson *et al.*, 2016).

Accommodation issues and family support

The high rates of precarious accommodation in this population imply there is a need for better family support for TGD young people. We inferred that, because of the age range of our participants, family support will often form a vital component of stable accommodation. A high proportion of participants (65.8%) reported that they lacked family support, and this was associated with poor mental health outcomes. Research has shown that trans young people who are supported in their identity by their parents have fewer difficulties with mental health, are less likely to report

suicidal ideation and are much more likely to have a secure/stable home to live in (Travers *et al.*, 2012).

Educational environments (school, university and technical college)

Most participants in this study reported negative experiences within education environments, and these were significantly associated with all adverse health outcomes measured. This is in line with previous research that has established that LGBTIQ young people are not adequately supported in school settings, and that this lack of support is associated with negative health outcomes (Jones, 2015; Bradlow *et al.*, 2017). TGD young people in schools who are not supported by teachers are more likely to experience abuse at school and are more likely to have worse educational outcomes than gender diverse young people who do feel supported by their teachers; e.g. through the use of correct language (Jones *et al.*, 2016). Programs that proactively and equitably support TGD young people in educational settings are necessary to mitigate these transphobic experiences.

Bullying and discrimination

The high rates of bullying (74%) and discrimination (68.9%) reported by participants underscore the need for broader interventions that target public perceptions through promoting acceptance and understanding of gender diversity. Gender diverse young people in Australia are more likely to be exposed to homophobic and/or transphobic abuse, including physical abuse, than their same-gender attracted (LGB) peers (Jones and Hillier, 2013). These findings highlight the need for anti-discrimination and anti-bullying programs that are specific to gender diverse young people.

Implications

These results show that TGD young people are a marginalised group that urgently needs interventions specifically targeted to improve their mental health. The findings show the need for improved protections for TGD young people to reduce many of the factors that are associated with poor mental health, including policies to decrease discrimination, bullying, abuse and other negative experiences that TGD young people are exposed to. These policies should be considered for implementation in educational, clinical and support settings. There is also a need to improve general understanding and acceptance in the general public. Given these high rates of mental health difficulties, it is also vital that TGD young people are able to access effective, safe and TGD-friendly mental health care providers.


Study limitations

The study design was cross-sectional, and therefore causal pathways cannot be inferred. Longitudinal data are necessary to more fully investigate the temporal relationships between adverse life events and mental health outcomes. In addition, we surveyed only Australian TGD young people, and results may not be generalisable to other countries, although our findings are remarkably consistent with international literature. There is potential for self-selection bias due to the survey's online nature. In addition, people without internet access were automatically excluded from the survey. The majority of participants were assigned female at birth, and therefore these data may not be representative of the wider TGD population. Further research specific to Aboriginal and

Torres Strait Islander TGD young person populations should be developed in collaboration with Aboriginal and Torres Strait Islander communities.

Conclusion

These findings support previous international research indicating that TGD populations experience mental health issues at higher rates than cisgender populations. It is significant that these rates of mental health issues are being reported in a community-based sample of young people, rather than a clinical sample. This study also demonstrates that TGD young people report alarmingly high rates of negative experiences, including discrimination and bullying. These negative and transphobic experiences are associated with poor mental health of TGD young people. Young people need to be supported by their peers, families, school and work peers to achieve optimal levels of mental wellbeing. Furthermore, services – including schools – need to ensure that they are gender inclusive to respond effectively and appropriately to the mental health needs of TGD young people outlined in this paper. These measures should be taken proactively as preventative measures, rather than as reactive measures, to create equitable spaces for all young people where they can thrive. This would help to prevent poor mental health in this population. As more TGD young people seek support from medical and mental health services in Australia and worldwide, it is crucial that clinical service providers are aware of the mental health issues faced by gender diverse young people and that they can offer TGD competent health care.

Author ORCIDs.  Penelope Strauss, 0000-0002-4828-0983.

Financial support. This research received no specific grant from any funding agency, commercial or not-for-profit sectors. PS is supported by an Australian Government Research Training Program Scholarship at the University of Western Australia. AL is supported by an NHMRC Career Development Fellowship (#1148793).

Conflict of interest. None.

Ethical standards. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

References

- Adams N, Hitomi M and Moody C (2017) Varied reports of adult transgender suicidality: synthesizing and describing the peer-reviewed and gray literature. *Transgender Health* 2, 60–75.
- Australian Bureau of Statistics (2007) *National Survey of Mental Health and Wellbeing: Summary of Results*. cat. no. 4326.0. ABS: Canberra, Australia.
- Australian Bureau of Statistics (2014) *Estimates and Projections, Aboriginal and Torres Strait Islander Australians, 2001 to 2026*. cat. no. 3238.0. ABS: Canberra, Australia.
- Australian Bureau of Statistics (2016) *Australian Demographic Statistics, Dec 2015*. cat. no. 3101.0. ABS: Canberra, Australia.
- Bouman WP, Claes L, Brewin N, Crawford JR, Millet N, Fernandez-Aranda F and Arcelus J (2017) Transgender and anxiety: a comparative study between transgender people and the general population. *International Journal of Transgenderism* 18, 16–26.
- Bradlow J, Bartram F, Guasp A and Jadv V (2017) *School Report: The Experiences of Lesbian, Gay, Bi and Trans Young People in Britain's Schools in 2017*. Stonewall; Centre for Family Research, University of Cambridge.

- Cohen-Kettenis PT, Steensma TD and de Vries ALC (2011) Treatment of adolescents with gender dysphoria in the Netherlands. *Child and Adolescent Psychiatric Clinics of North America* **20**, 689–700.
- Collier K, van Beusekom G, Bos H and Sandfort T (2013) Sexual orientation and gender identity/expression related peer victimization in adolescence: a systematic review of associated psychosocial and health outcomes. *Journal of Sex Research* **50**, 299–317.
- Flores AR, Herman JL, Brown TNT and Conron KJ (2017) *Age of Individuals Who Identify as Transgender in the United States*. Los Angeles, CA: The Williams Institute.
- Gordon AR, Conron KJ, Calzo JP, White MT, Reisner SL and Austin SB (2018) Gender expression, violence, and bullying victimization: findings from probability samples of high school students in 4 US school districts. *Journal of School Health* **88**, 306–314.
- Hillier L, Jones T, Monagle M, Overton N, Gahan L, Blackman J and Mitchell A (2010) *Writing Themselves in 3 (WTi3)*. Melbourne, Australia: Australian Research Centre in Sex, Health and Society, La Trobe University.
- Hyde Z, Doherty M, Tilley PJM, McCaul KA, Rooney R and Jancey J (2014) *The First Australian National Trans Mental Health Study: Summary of Results*. Perth, Australia: School of Public Health, Curtin University.
- Inch E (2016) Changing minds: the psycho-pathologization of trans people. *International Journal of Mental Health* **45**, 193–204.
- Jones T (2015) *Policy and Gay, Lesbian, Bisexual, Transgender and Intersex Students*. Cham: Springer International Publishing.
- Jones T and Hillier L (2013) Comparing trans-spectrum and same-sex-attracted youth in Australia: increased risks, increased activism. *Journal of LGBT Youth* **10**, 287–307.
- Jones T, Smith E, Ward R, Dixon J, Hillier L and Mitchell A (2016) School experiences of transgender and gender diverse students in Australia. *Journal of Sex Education* **16**, 1–16.
- Kroenke K, Spitzer R and Williams J (2001) The PHQ-9: validity of a brief depression severity measure. *Journal of General Internal Medicine* **16**, 606–613.
- Lawrence D, Johnson S, Hafekost J, Boterhoven De Haan K, Sawyer M, Ainley J and Zubrick S (2015) *The Mental Health of Children and Adolescents. Report on the Second Australian Child and Adolescent Survey of Mental Health and Wellbeing*. Australian Government: Canberra, Australia.
- McDermott E, Hughes E and Rawlings V (2017) The social determinants of lesbian, gay, bisexual and transgender youth suicidality in England: a mixed methods study. *Journal of Public Health* **40**, e244–e251.
- Mustanski B and Liu R (2013) A longitudinal study of predictors of suicide attempts among lesbian, gay, bisexual, and transgender youth. *Archives of Sexual Behavior* **42**, 437–448.
- Olson KR, Durwood L, Demeules M and McLaughlin KA (2016) Mental health of transgender children who are supported in their identities. *Pediatrics* **137**, 1–8.
- Reisner SL, Veters R, Leclerc M, Zaslow S, Wolfrum S, Shumer D and Mimiaga MJ (2015) Mental health of transgender youth in care at an adolescent urban community health center: a Matched Retrospective Cohort Study. *Journal of Adolescent Health* **56**, 274–279.
- Reisner SL, Poteat T, Keatley J, Cabral M, Mothopeng T, Dunham E, Holland CE, Max R and Baral SD (2016) Global health burden and needs of transgender populations: a review. *The Lancet* **338**, 412–436.
- Rider GN, McMorris BJ, Gower AL, Coleman E and Eisenberg ME (2018) Health and care utilization of transgender and gender nonconforming youth: a population-based study. *Pediatrics* **141**, e20171683.
- Roberts A, Rosario M, Slopen N, Calzo JP and Bryn Austin S (2013) Childhood gender nonconformity, bullying victimization, and depressive symptoms across adolescence and early adulthood: an 11-year longitudinal study. *Journal of the American Academy of Child & Adolescent Psychiatry* **52**, 143–152.
- Rood B, Reisner S, Surace F, Puckett J, Maroney M and Pantalone D (2016) Expecting rejection: understanding the minority stress experiences of transgender and gender-nonconforming individuals. *Transgender Health* **1**, 151–164.
- Smith E, Jones T, Ward R, Dixon J, Mitchell A and Hillier L (2014) *From Blues to Rainbows: Mental Health and Wellbeing of Gender Diverse and Transgender Young People in Australia*. Melbourne, Australia: The Australian Research Centre in Sex, Health and Society.
- Spitzer R, Kroenke K, Williams J and Lowe B (2006) A brief measure for assessing generalized anxiety disorder – The GAD-7. *Archives of Internal Medicine* **166**, 1092–1097.
- Taliaferro LA, McMorris BJ, Rider GN and Eisenberg ME (2018) Risk and protective factors for self-harm in a population-based sample of transgender youth. *Archives of Suicide Research* **23**, 203–221.
- Telfer M, Tollit M and Feldman D (2015) Transformation of health-care and legal systems for the transgender population: the need for change in Australia. *Journal of Paediatrics and Child Health* **51**, 1051–1053.
- Testa RJ, Hendricks ML, Goldblum P and Bradford J (2012) Effects of violence on transgender people. *Professional Psychology: Research and Practice* **43**, 452–459.
- Testa RJ, Michaels MS, Bliss W, Rogers ML, Balsam KF and Joiner T (2017) Suicidal ideation in transgender people: gender minority stress and interpersonal theory factors. *Journal of Abnormal Psychology* **126**, 125–136.
- Travers R, Bauer G, Pyne J, Bradley K, Gale L and Papadimitriou M (2012) *Impacts of Strong Parental Support for Trans Youth*. Children's Aid Society of Toronto & Delisle Youth Services: Toronto.
- Veale JF, Watson RJ, Peter T and Saewyc EM (2017) Mental health disparities among Canadian transgender youth. *Journal of Adolescent Health* **60**, 44–49.
- Vrouenraets LJ, Fredriks AM, Hannema SE, Cohen-Kettenis PT and de Vries MC (2015) Early medical treatment of children and adolescents with gender dysphoria: an empirical ethical study. *Journal of Adolescent Health* **57**, 367–373.
- Winter S, Diamond M, Green J, Karasic D, Reed T, Whittle S and Wylie K (2016) Transgender people: health at the margins of society. *The Lancet* **388**, 390–400.