IAC Ch 7, p.1

641—7.4(139A) Required immunizations.

7.4(1) Applicants enrolled or attempting to enroll shall have received the following vaccines in accordance with the doses and age requirements listed below:

IMMUNIZATION REQUIREMENTS

Institution	Age	Vaccine	Total Doses Required
Licensed child care center	< 2 months	None	
	2 through 5 months	Diphtheria/Tetanus/Pertussis ¹ Polio <i>Haemophilus influenzae</i> type B	1 dose 1 dose 1 dose
	6 through 14 months	Diphtheria/Tetanus/Pertussis ¹ Polio <i>Haemophilus influenzae</i> type B	2 doses 2 doses 2 doses
	15 through 18 months	Diphtheria/Tetanus/Pertussis ¹ Polio <i>Haemophilus influenzae</i> type B	3 doses 3 doses 3 doses, with the final dose in the series received when the applicant is ≥ 12 months of age; or 1 dose received when the applicant is ≥ 15 months of age.
		Measles/Rubella ²	1 dose of measles/rubella- containing vaccine received after the applicant was at least 12 months of age; or the applicant demonstrates a positive antibody test for measles and rubella.
	19 months and older	Diphtheria/Tetanus/Pertussis ¹ Polio Haemophilus influenzae type B	3 doses 3 doses 3 doses, with the final dose in the series received when the applicant is ≥ 12 months of age; or 1 dose received when the applicant is ≥ 15 months of age.
		Measles/Rubella ²	1 dose of measles/rubella- containing vaccine received after the applicant was at least 12 months of age; or the applicant demonstrates a positive antibody test for measles and rubella.
		Varicella	1 dose if the applicant was born on or after September 15, 1997, unless the applicant has had a reliable history of natural disease. This dose shall have been received after the applicant was at least 12 months of age.

Ch 7, p.2

Institution	Age	Vaccine	Total Doses Required
Elementary school/ secondary school	4 years of age and older	Diphtheria/Tetanus/Pertussis ^{1,3}	3 doses, with at least 1 dose of diptheria/tetanus/pertussis-containing vaccine received after the applicant's fourth birthday if the applicant was born on or before September 15, 2000; or 4 doses, with at least 1 dose of diptheria/tetanus/ pertussis-containing vaccine received after the applicant's fourth birthday if the applicant was born after September 15, 2000. Applicants ≥ 7 years of age are exempt from receiving further doses of pertussis-containing vaccine; therefore, tetanus and diphtheria-containing vaccine should be used.
		Polio	3 doses, with at least 1 dose on or after the applicant's fourth birthday.
		Measles/Rubella ²	2 doses of measles/rubella- containing vaccine; or the applicant demonstrates a positive antibody test for measles and rubella. The first dose shall have been received on or after the applicant's first birthday; the second dose shall have been received no less than 28 days after the first dose.
		Hepatitis B	3 doses if the applicant was born on or after July 1, 1994.
		Varicella	1 dose if the applicant was born on or after September 15, 1997, unless the applicant has had a reliable history of natural disease. This dose shall have been received after the applicant was at least 12 months of age.

¹ Pediatric diphtheria and tetanus vaccine may be substituted for the pertussis-containing vaccine, without a medical exemption, when pertussis vaccine is contraindicated for the child < 7 years of age.

² Mumps vaccine may be included in measles/rubella-containing vaccine.

³ If a child received the first dose of tetanus/diphtheria-containing product when the child was < 12 months of age, 4 doses are required, with 1 dose on or after the child's fourth birthday. If a child received the first dose of tetanus/diphtheria-containing product when the child was ≥ 12 months of age, 3 doses are required, with 1 dose on or after the child's fourth birthday.

^{7.4(2)} Vaccine doses administered ≤ 4 days before the minimum interval or age shall be counted as valid. Doses administered ≥ 5 days earlier than the minimum interval or age shall not be counted as valid doses and shall be repeated as age-appropriate.

^{7.4(3)} For vaccine administration, the minimum age and intervals recommended by the advisory committee on immunization practices shall be followed.