



December 5, 2017

The Honorable Kim Reynolds and
Iowa General Assembly
State Capitol
1007 E. Grand Avenue
Des Moines, Iowa 50319

Dear Governor Reynolds and Iowa General Assembly:

I am pleased to submit to you the Department's *2016 Iowa Statewide Greenhouse Gas Emissions Inventory Report and Technical Support Document* as required by Iowa Code 455B.104. This policy-neutral report is eleventh statewide greenhouse gas inventory conducted for Iowa.

We hope this report is useful for policy makers as they continue to discuss this important issue.

Sincerely,

A handwritten signature in black ink that reads "Chuck Gipp".

Chuck Gipp
Director
Department of Natural Resources

With Enclosures



IOWA DEPARTMENT OF NATURAL RESOURCES

2016 Iowa Statewide Greenhouse Gas Emissions Inventory Report

Required by Iowa Code 455B.104

December 5, 2017

Iowa Department of Natural Resources
502 E. 9th Street
Des Moines, IA 50319

Table of Contents

Background 3

2016 Statewide GHG Emissions 3

GHG Emissions by Sector 7

GHG Emissions by Pollutant..... 12

Comparison with U.S. Emissions..... 14

Future Emissions 15

Future Improvements 17

Background

This report is required by Iowa Code 455B.104, which requires the Iowa DNR to estimate greenhouse gas (GHG) emissions during the previous year and forecast trends in emissions. The report must be submitted to the Governor and Iowa General Assembly by December 31 each year. This report focuses on calendar year 2016 GHG emissions and includes emissions of six GHGs: carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), perfluorocarbons (PFC), hydrofluorocarbons (HFC) and sulfur hexafluoride (SF₆).

This inventory is based on statewide activity data from the following sectors:

- agriculture
- fossil fuel combustion
- industrial processes
- natural gas transmission and distribution
- transportation
- solid waste
- wastewater treatment
- land use, land use change, and forestry (LULUCF)

Emissions were calculated using the U.S. Environmental Protection Agency's (EPA) State Inventory Tool (SIT) and self-reported emissions data from landfills, industrial facilities and power plants. The calculation method and uncertainty for each sector are discussed in detail in the Iowa DNR's Technical Support document available on the DNR's [Greenhouse Gas Emissions Inventory webpage](#). Benefits of preparing this report include the ability to evaluate emissions trends and develop a baseline to track progress in reducing emissions. A state-specific inventory also provides a more in-depth analysis and more accurate inventory of emissions compared to national emissions.

2016 Statewide GHG Emissions

In 2016, total gross Iowa greenhouse gas emissions were 128.28 million metric tons carbon dioxide equivalents (MMtCO₂e) as shown in Table 1. This is a decrease of 2.61 MMtCO₂e (2.00%) from 2015. Most of this decrease comes from a 4.13 MMtCO₂e reduction from power plants due to mild weather, decreased electric generation from coal (-14.12%) and increased generation from zero-emitting sources such as wind (12.18%) and solar.¹

These decreases helped to offset increases in emissions from industrial processes (+4.28%), LULUCF (+4.69%), agriculture (+1.82%) and other sectors. Overall, total 2016 statewide GHG emissions were 9.29% lower than their peak in 2007 as shown in Figures 1 and 2.

¹ Solar generation in 2015 rounded to zero million megawatt hours. Solar generation in 2016 was 416 million megawatt hours.

Table 1: GHG Emissions 2005 – 2016 by Sector ((Million Metric Tons Carbon Dioxide Equivalents (MMtCO₂e))²)

Emissions (MMtCO ₂ e)	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	Change from 2015		
													MMtCO ₂ e	%	
Agriculture	33.60	35.71	40.23	36.65	36.48	36.38	37.91	36.67	37.03	37.76	38.78	39.48	+0.71	+1.82%	↑
Power Plants	36.83	36.35	40.04	41.78	37.71	42.33	38.98	35.76	33.06	33.44	29.46	25.33	-4.13	-14.02%	↓
Residential, Commercial, and Industrial Fuel Use	27.03	27.26	28.55	31.34	30.73	31.23	31.44	29.96	32.82	32.82	31.54	31.90	+0.36	+1.16%	↑
Industrial Processes	4.57	4.71	4.78	4.92	4.22	4.79	4.49	5.17	5.34	5.28	5.28	5.50	+0.23	+4.28%	↑
Land Use, Land Use Change, and Forestry (LULUCF) ³			3.40				0.67	0.48		3.27	2.99	3.13	+0.14	+4.69%	↑
Natural Gas Transmission and Distribution	1.37	1.37	1.39	1.40	1.40	1.39	1.40	1.40	1.40	1.40	1.40	1.41	+0.01	+0.62%	↑
Transportation	20.14	20.35	19.96	19.99	19.40	19.41	19.58	19.59	19.46	19.55	19.31	19.36	+0.05	+0.28%	↑
Waste	3.10	3.03	3.07	3.10	3.06	2.01	1.94	2.15	1.96	1.94	2.14	2.16	+0.02	+1.00%	↑
Total Gross Emissions	126.63	128.77	141.42	139.16	133.00	137.54	136.39	131.18	131.07	135.46	130.90	128.28	-2.61	-2.00%	↓
Carbon Stored in LULUCF ⁴	-20.56	-5.81	0	-3.92	-5.00	-2.01	0	0	-0.71	0	0	0			
Total Net Emissions	106.07	122.96	141.42	135.25	128.00	135.54	136.39	131.18	130.36	135.46	130.90	128.28	-2.61	-2.00%	↓

² Totals may not equal the exact sum of subtotals in this table due to independent rounding Values may not match values in the previous 2015 inventory published by the Department in December 2016 as they have been recalculated using the IPCC AR4 GWP. The adjustments are described in detail in the Technical Support Document.

³ Carbon is emitted in some years from the LULUCF sector, but carbon is stored in the LULUCF sector in other years. Emissions from the LULUCF sector are shown in this row as positive numbers.

⁴ Carbon stored in LULUCF is shown in this row as a negative number.

Figure 1: Iowa Gross GHG Emissions 2005 – 2016 (MMtCO₂e)





Figure 2: Iowa Gross⁵ GHG Emissions 2005 – 2016 by Sector (MMtCO₂e)




⁵ Does not include carbon storage or emissions from land use, land use change, and forestry (LULUCF).

GHG Emissions by Sector

The majority of GHG emissions in Iowa in 2016 were from the agriculture sector and from fossil fuel use by power plants and the residential/commercial/ industrial (RCI) sectors. Together the emissions from power plants and RCI fuel use account for nearly half (44.61%) of the state's GHG emissions as shown in Figure 3.

Figure 3: 2016 Iowa GHG Emissions by Sector⁶



⁶ Industrial fossil fuel use refers to GHG emissions from fossil fuels combusted by industrial facilities. Industrial processes means GHGs emitted during the production of or use of specific products such as ammonia, urea, nitrogen, cement, iron, steel, lime, etc.

Emissions from each sector are summarized below. Please refer to the [2016 GHG Inventory Technical Support Document](#) for more information on a specific sector, such as sources of emissions, calculations and uncertainty.

Agriculture

This sector includes GHG emissions from livestock and crop production such as enteric fermentation, manure management and agricultural soils. Enteric fermentation includes emissions from the digestive systems of ruminant animals. Emissions from agricultural soils include emissions from manure, runoff, plant fertilizers, plant residues and cultivation of highly organic soils. GHG emissions from fossil-fuel fired agricultural equipment (such as tractors) are included in the transportation sector. As shown in Table 2, agriculture emissions increased 1.82% from the previous year due to increases in both livestock and crop production.

Table 2: GHG Emissions from Agriculture (MMtCO₂e)

Category	2015	2016	% Change
Enteric Fermentation	8.31	8.38	+0.83%
Manure Management	10.04	9.86	-1.74%
Agricultural Soil Management	20.43	21.24	+3.97%
Total	38.78	39.48	+1.82%

Fossil Fuel Combustion

This sector includes GHG emissions from fossil fuels combusted in four categories: power plants, residential, industrial and commercial (The residential, industrial and commercial categories are often combined into one category called RCI). Together, these four categories account for 44.61% of Iowa's GHG total emissions. Emissions from power plants decreased by 4.13 MMtCO₂e (14.02%) from the previous year due to decreased combustion of coal and increases in the amount of electricity generated from lower-emitting fuels, such as natural gas, and zero-emitting fuels, such as wind and solar. Emissions from the other three fossil fuel combustion categories increased as shown in Table 3.

Table 3: GHG Emissions from Fossil Fuel Combustion (MMtCO₂e)

Category	2015	2016	% Change
Power Plants	29.56	25.33	-14.02%
Residential, Commercial, Industrial (RCI)	31.54	31.90	+1.16%
Residential	4.49	4.55	+1.27%
Commercial	4.60	4.76	+3.52%
Industrial	22.44	22.59	+0.65%
Total	61.00	57.23	-6.18%

Residential, Commercial, Industrial (RCI)


Actual fuel use data for 2016 for the RCI sector was not available from the U.S. Energy Information Administration (EIA), so emissions were calculated based on projected energy consumption values from the EIA's *Annual Energy Outlook 2017 with Projections to 2040*. Emissions predicted for 2015 from the RCI sector in last year's inventory (30.96 MMtCO₂e) were replaced with actual 2015

consumption values now available from EIA. The resulting recalculated 2015 emissions were 31.54 MMTCO₂e. Emissions from this sector increased because the consumption of fossil fuels by users in each category increased.

Power Plants

This category includes emissions from fossil fuels that are combusted at power plants to generate electricity. The DNR used emissions reported by power plants to EPA as required by the federal GHG reporting program (40 Code of Federal Regulations 98). The CO₂ emissions for these facilities are measured by continuous emission monitors (CEMS). Emissions from this sector decreased as electric generation from coal decreased and generation from renewables and natural gas increased as shown in Figure 4. Combustion of natural gas emits approximately half as much CO₂ as is emitted from the combustion of coal. CO₂ is not emitted when electricity is generated from “zero-emitting carbon sources” such as wind, hydroelectric or nuclear power.

Figure 4: 2015 and 2016 Net Iowa Electric Generation by Energy Source (Million Megawatt Hours)⁷



Industrial Processes

This sector includes non-combustion GHG emissions from a variety of processes including cement production, lime manufacture, limestone and dolomite use, soda ash use, iron and steel production, ammonia production, nitric acid production, substitutes for ozone depleting substances (ODS) and electric power transmission and distribution. GHG emission trends in each process category vary, but overall total industrial process emissions increased 4.28% from 2015 - 2016 as shown in Table 4.

⁷ U.S. EIA: [Net Generation by State by Type of Producer by Energy Source](#), September 2017.

GHG emissions reported by industrial facilities to EPA as required by the federal GHG reporting program were used for these categories: ammonia and urea production, cement manufacture, iron and steel production, lime manufacture and nitric acid production. Emissions from the other categories were calculated using EPA's SIT.

Table 4: GHG Emissions from Industrial Processes (MMtCO₂e)

Category	2015	2016	% Change
Ammonia and Urea Production	0.81	0.92	+13.59%
Cement Manufacture	1.50	1.58	+4.88%
Electric Power Transmission & Distribution Systems	0.05	0.05	NA ⁸
Iron and Steel Production	0.16	0.19	+14.80%
Lime Manufacture	0.13	0.15	+9.73%
Limestone and Dolomite Use	0.21	0.21	NA ⁸
Nitric Acid Production	0.74	0.75	+1.07%
Ozone Depleting Substance Substitutes	1.64	1.63	-0.19%
Soda Ash Consumption	0.02	0.02	-2.48%
Total	5.28	5.50	4.28%

Natural Gas Transmission and Distribution (T & D)

This sector includes emissions from natural gas transmission and distribution systems in the state. GHG emissions increased 0.62% from 2015 as shown in Table 5, due to an increase in the number entities/customers connected to natural gas service in the state.

Table 5: GHG Emissions from Natural Gas Transmission and Distribution (MMtCO₂e)⁹

Category	2015	2016	% Change
Transmission	0.7868	0.7867	-0.01%
Distribution	0.6168	0.6205	+1.42%
Total	1.4031	1.4073	+0.62%

Transportation

The transportation sector includes GHG emissions from both highway and non-highway vehicles. Aviation, boats, locomotives, tractors, other utility vehicles and alternative fuel vehicles are considered non-highway vehicles. Emissions from highway vehicles are calculated based on vehicle miles traveled, while emissions from non-highway vehicles are calculated based on fuel consumption. Total vehicle miles traveled by lowans increased 0.47%, resulting in a 0.28% increase in GHG emissions as shown in Table 6.

Table 6: GHG Emissions from Transportation (MMtCO₂e)

Category	2015	2016	% Change
Transportation	19.31	19.36	+0.28%

⁸ Due to lack of current data, the DNR assumed 2015 emissions = 2016 emissions.

⁹ DNR uses two decimal places throughout this report for consistency. However, in this sector four decimal places are needed to show the difference in emissions from year to year.

Waste

The waste sector includes GHG emissions from both solid waste landfills and the treatment of municipal and industrial wastewater. Overall, GHG emissions from waste increased 1.00% from 2015 as shown in Table 7 due to more solid waste being placed in landfills. DNR primarily used facility-specific emissions data directly reported by facilities to U.S. EPA. Emissions from smaller landfills that are not required to report to EPA were estimated using the LandGEM model. The SIT was used to estimate emissions from municipal wastewater treatment and combustion of municipal solid waste.

Table 7: GHG Emissions from Waste (MMtCO₂e)

Category	2015	2016	% Change
Solid Waste	1.69	1.71	+1.17%
Wastewater	0.45	0.45	+0.35%
Total	2.14	2.16	+1.00%

Land Use, Land Use Change, and Forestry (LULUCF)

The LULUCF sector includes emissions from liming agricultural soils and fertilizing lawns, golf courses and other landscaping (settlement soils). It also includes carbon sequestered by forests and urban trees, as well as carbon stored in yard waste and food scraps that are sent to the landfill. Data that is more current was used to recalculate carbon sequestration from forests and trees in 2015, and the 2015 value was used as a surrogate for 2016. This recalculation is discussed in detail in the Technical Support Document.

Overall, total 2016 emissions from LULUCF were 3.13 MMtCO₂e as shown in Table 8. This is a 4.69% increase in the CO₂e being emitted. This can be attributed to increases in emissions from liming agricultural soils and fertilizing settlement soils (e.g. landscaping, lawns, golf courses). Due to a lack of current data, carbon sequestration from forest carbon flux and urban trees was assumed unchanged from 2015, as were emissions from urea fertilization and sequestration from yard trimmings and food scraps in landfills. Emissions from forest fires were not calculated due to a lack of data.

Table 8: GHG Emissions by LULUCF (MMtCO₂e)

Category	2015	2016	% Change
Forest Carbon Flux	2.87	2.87	NA ¹⁰
Liming of Agricultural Soils	0.34	0.46	+37.22%
Urea Fertilization	0.15	0.15	NA ¹⁰
Urban Trees	-0.74	-0.74	NA ¹⁰
Yard Trimmings & Food Scraps in Landfills	-0.12	-0.12	NA ¹⁰
Fertilization of Settlement Soils	0.49	0.51	+3.12%
Total	2.99	3.13	+4.69%

Carbon emitted or sequestered from agricultural soil carbon flux are not included in the inventory. This is because the SIT does not have a calculation function for this category and because of the uncertainty in this category. Scientific studies and literature reviews do not agree on the relationship between soil

¹⁰ Due to a lack of current data, the DNR assumed 2015 values = 2016 values.


tillage and soil carbon. Therefore, the DNR did not include this category. More details on the uncertainty in soil carbon flux are included in the [2016 GHG Inventory Technical Support Document](#).

GHG Emissions by Pollutant

GHGs included in the inventory are carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), perfluorocarbons (PFC), hydrofluorocarbons (HFC) and sulfur hexafluoride (SF₆). Figures 5-8 show the distribution of GHG pollutants in 2016 by both pollutant and by category.

Carbon dioxide (CO₂) is the greenhouse gas emitted in the highest amounts in Iowa, accounting for 66.37% of all greenhouse gas emissions in 2016. Nearly all CO₂ emissions are from fossil fuel combustion and transportation as shown in Figure 5, with a small percentages coming from industrial processes such as the production of cement, lime, ammonia, urea, iron and steel; as well as the use of limestone, dolomite and soda ash in manufacturing.

Figure 5: 2016 Carbon Dioxide Emissions by Sector



Methane (CH₄) and nitrous oxide (N₂O) were emitted in smaller amounts, and the majority of these two pollutants are from agriculture as shown in Figures 6 and 7. CH₄ emissions were 20.86 MMtCO₂e or 16.26% of total 2016 GHG emissions. N₂O emissions in 2016 were 24.94 MMtCO₂e or 18.66% of total GHG emissions.

Figure 6: 2016 Methane Emissions by Sector






Figure 7: 2016 Nitrous Oxide Emissions by Sector



Emissions of Hydrofluorocarbons (HFCs), perfluorocarbons (PFCs) and sulfur hexafluoride (SF₆) are accounted for in the Industrial Processes sector. They are emitted either from substitutes for ODS or as insulation in electric power transmission and distribution lines as shown in Figure 8. In 2016, emissions of these three pollutants totaled 1.68 MMtCO₂e or 1.31% 2016 Iowa GHG emissions.

Figure 8: 2016 HFC/PFC/SF₆ Emissions by Sector



Comparison with U.S. Emissions

Figures 9 and 10 compare Iowa and national GHG emissions by sector. Overall, Iowa emitted 1.95% of total 2015 U.S. GHG emissions. Agricultural emissions account for a greater percentage of GHG emissions in Iowa than in the total U.S., which is logical given Iowa's substantial agricultural economy.

The fossil fuel combustion, natural gas distribution and transmission, and transportation sectors have been combined into one sector called "Energy" to be consistent with the national GHG inventory. Emissions from 2015 are used for this comparison, as the 2016 national GHG inventory has not yet been published.

Figure 9: 2015 U.S. GHG Emissions by Sector





Figure 10: 2015 Iowa GHG Emissions by Sector



Future Emissions

Iowa Code 455B.104 requires that the DNR forecast trends in GHG emissions. The DNR projected emissions from 2016 to 2030 using the SIT Projection Tool. As with many forecasts, numerous factors affect the level of uncertainty with future emissions. These factors may include among other things – the economy, weather, current and future environmental regulations, energy efficiency and conservation practices, driving practices, use of renewable fuels, etc.

The SIT Projection Tool forecasts emissions from industrial processes, agriculture and waste based on historical emissions from 1990 – 2014, using a combination of data sources and national projections for activity data. The average annual change in emissions that the SIT projects are shown in Table 9.

Table 9: Average Annual Projected Change in Emissions 2017 - 2030

Sector	% Ave. Annual Change
Agriculture	+0.17%
Electric Power Generation Fossil Fuel Use	-2.05%
RCI Fuel Use	+0.27%
Industrial Processes	+3.48%
Natural Gas Transmission & Distribution	+0.71%
Transportation	-0.90%
Waste	+0.98%
Total Emissions	-0.24%

Short-term Projections for the Electric Power Sector

In April 2017, the U.S. Energy Information Administration of the Department of Energy announced that CO₂ emissions in the national energy sector fell 1.7% from 2015 – 2016 due to a reduction in electricity generated from coal and an increase in generation from natural gas and renewable energy. Weather also played a role, as warmer years may result in reduced energy consumption because it typically costs less to cool a home than heat it. Heating days in the U.S. in 2016 were the second lowest of any year in the past 67 years.¹¹

However, the most recent emissions data available for Iowa power plants does not follow the same trend as national emissions. Data from EPA's Clean Air Markets Division shows that CO₂ emissions from the electric power generation during the first six months of 2017 are 15.05% higher than CO₂ emissions from the first six months of 2016. It is unclear what caused this trend, but some factors may be a change in the number of heating and/or cooling days, changes in the prices of different fuels or other market forces.

Although the SIT Projection Tool provides a good first look at projected future emissions, it has several areas of uncertainty:

1. Agricultural emissions are highly dependent on the weather and crop and livestock prices, which are not addressed by the Projection Tool.

¹¹ U.S. EIA: [Today in Energy - April 10, 2017](#).

2. In sectors where the Projection Tool predicts future emissions based on historical emissions, it only uses emissions from 1990 – 2014 and does not consider 2015 - 2016 emissions.
3. The Projection Tool forecasts emissions from fossil fuel use based on the reference case from the EIA's *Annual Energy Outlook (AEO) 2016 with Projections to 2040*, which projects emissions at the regional level and not the state level. A more current reference case, *AEO 2017*, has been published but is not used by the tool.
4. The *AEO 2016* used includes implementation of the CPP, which is currently stayed and EPA proposes to repeal.
5. The Projection Tool does not address publicly announced changes to Iowa's fossil fuel generation mix:
 - Iowa utilities have announced that from 2016 - 2025, approximately 1,000 MW of coal-fired electric generation units will retire or convert to natural gas. During that same period, approximately 185 MW of older natural gas-fired electric generation units will retire, and approximately 650 MW of newer, more efficient natural gas-fired electric generating units will come online. This will significantly reduce emissions from the electric power sector as natural gas emits approximately 50% less CO₂ per heating unit than coal emits.
 - At least 3,200 MW of additional renewables are planned for 2017 - 2020.

Future Improvements

The DNR continually strives to make the annual statewide GHG inventory as accurate and timely as possible. Possible areas for enhancement include improved forecasting and a more extensive literature review of research regarding soil carbon. Additionally, the DNR plans to work with U.S. EPA GHG inventory staff to investigate whether emissions from the application of synthetic fertilizer are double-counted in the agriculture and LULUCF sectors.



IOWA DEPARTMENT OF
NATURAL RESOURCES

2016 Iowa Statewide
Greenhouse Gas Emissions
Inventory Report

Technical Support Document

Required by Iowa Code 455B.104

December 5, 2017

Iowa Department of Natural Resources
502 E. 9th Street
Des Moines, IA 50319

This page is left intentionally blank.

Acronyms and Key Terms.....	4
Chapter 1 – General Calculation Method	6
Chapter 2 - Agriculture.....	9
Chapter 3 – Fossil Fuel Consumption.....	20
Chapter 4 - Industrial Processes	26
Chapter 5 - Natural Gas Transmission & Distribution.....	31
Chapter 6 - Transportation	34
Chapter 7 – Waste: Solid Waste	39
Chapter 8 – Waste: Wastewater Treatment.....	42
Chapter 9 - Land Use, Land Use Change, and Forestry (LULUCF)	46
Chapter 10 – Electricity Consumption	51
Forecasting.....	54
References	54
Appendix A – Iowa GHG Emissions 2005 – 2016 by Sector	63
Appendix B – Iowa GHG Emissions 2005 – 2016 by Pollutant.....	65

Acronyms and Key Terms

AEO	Annual Energy Outlook
AR4	Fourth Assessment Report
BOD	biochemical oxygen demand
BOF	blast oven furnace
Btu	British thermal unit
CAMD	Clean Air Markets Division
CH ₄	methane
CO ₂	carbon dioxide
COMET	Carbon Management and Evaluation Online Tool
CPP	Clean Power Plan
CRP	Conservation Reserve Program
DNR	Iowa Department of Natural Resources
DOT	United States Department of Transportation
EAF	electric arc furnace
EIA	United States Energy Information Administration
EIIP	Emission Inventory Improvement Program
EPA	United States Environmental Protection Agency
FERC	Federal Energy Regulatory Agency
FIDO	Forest Inventory Data Online
FHWA	Federal Highway Administration
FTA	Federal Transit Administration
GHG	greenhouse gas
GHGRP	Greenhouse Gas Reporting Program
HDGV	heavy duty gas vehicle
HDDV	heavy duty diesel vehicle
IDALS	Iowa Department of Agriculture and Land Stewardship
IDOT	Iowa Department of Transportation
IEA	International Energy Agency
ILPA	Iowa Limestone Producers Association
IPCC	Intergovernmental Panel on Climate Change
LDC	local distribution company
LDDT	light duty diesel truck
LDDV	light duty diesel vehicle
LDGT	light duty gasoline truck
LDGV	light duty gasoline vehicle
LFGTE	landfill gas to energy
LULUCF	land use, land use change, and forestry

Acronyms and Key Terms (Continued)

MC	motorcycle
MMtC	million metric tons carbon
MMtCO ₂ e	million metric tons carbon dioxide equivalent
MSW	municipal solid waste
N	nitrogen
NEI	National Emissions Inventory
NEMS	National Energy Modeling System
NRCS	Natural Resources and Conservation Service
NO ₃ ⁻	nitrates
NO ₂ ⁻	nitrites
N ₂ O	nitrous oxide
ODS	ozone depleting substance
OECD	Organization for Economic Co-operation and Development
PET	polyethylene terephthalate
PHMSA	Pipeline and Hazardous Materials Safety Administration
PS	polystyrene
PVC	polyvinyl chloride
SIT	State Inventory Tool
TAR	Third Assessment Report
USDA	United States Department of Agriculture
USFS	United States Forest Service
USGS	United States Geological Survey
VMT	vehicle miles traveled
WRI	World Resources Institute

Chapter 1 – General Calculation Method

Iowa Code 455B.104 requires that “by December 31 of each year, the department shall submit a report to the governor and the general assembly regarding the greenhouse gas (GHG) emissions in the state during the previous calendar year and forecasting trends in such emissions....” This Technical Support Document (TSD) provides documentation and additional calculations to support the [2016 Iowa Statewide Greenhouse Gas Emissions Inventory Report](#). Total Iowa GHG emissions from 2005 – 2016 are provided in Appendices A and B of this document. A state-specific inventory provides an in-depth analysis of emission trends and develops a baseline to track progress in reducing emissions

This inventory is based on statewide activity data from agriculture, fossil fuel combustion, industrial processes, natural gas transmission and distribution, transportation, solid waste, and wastewater treatment. It also includes carbon emitted or sequestered from land use, land use change, and forestry (LULUCF).

Method

Emissions were calculated using the most recent version of the United States Environmental Protection Agency’s (EPA) State Greenhouse Gas Inventory Tool (SIT)¹ and using available Iowa-specific activity data. The energy and industrial processes sectors were also supplemented with GHG emissions data submitted by individual Iowa facilities to the federal GHG reporting program (40 CFR 98).

The calculation methods in the SIT are based on the August 2004 version of EPA’s Emission Inventory Improvement Program (EIIP) guidance for greenhouse gases (ICF 2004). The individual modules for each sector are Excel workbooks populated with emission factors and default activity data for years 1990 – 2015, but allow the user to enter better state-specific activity data when it is available. Detailed information on the activity data used is provided in the corresponding chapter for each sector, under the “Method” heading. The individual modules then auto-calculate the resulting GHG emissions from each sector. The results from each module were then tabulated in an Excel spreadsheet. The SIT Projection Tool was then used to forecast emissions to 2030. The SIT modules and their corresponding chapters in this Technical Support Document are listed in Table 1. The coal module was not used, as there are no coal mines currently operating in Iowa.

¹ The SIT may be requested at <https://www.epa.gov/statelocalclimate/download-state-inventory-and-projection-tool>.

Table 1: TSD Chapters and Corresponding SIT Modules

TSD Chapter	SIT Module	Release Date	Pollutants Addressed
Agriculture	Ag	01/03/17	CH ₄ , N ₂ O
Energy	CO ₂ FFC	01/03/17	CO ₂
	Stationary Combustion	01/03/17	CH ₄ , N ₂ O
Industrial Processes	IP	01/03/17	CO ₂ , N ₂ O, HFC, PFC, SF ₆
Natural Gas Transmission and Distribution	Natural Gas and Oil	01/03/17	CH ₄
Transportation	Mobile Combustion	01/03/17	CO ₂ , CH ₄ , N ₂ O
Waste	Solid Waste	01/03/17	CO ₂ , CH ₄
	Wastewater	01/03/17	CH ₄ , N ₂ O
Land Use, Land Use Change, and Forestry (LULUCF)	LULUCF	01/03/17	CO ₂ , N ₂ O
Indirect Emissions from Electricity Consumption	Electricity Consumption	04/11/17	CO ₂
Future Emissions	Projection Tool	01/03/17	CO ₂ , CH ₄ , N ₂ O, HFC, PFC, SF ₆

Global Warming Potentials (GWP)

The potency of various greenhouse gases can vary, so greenhouse gas emissions are typically converted to a unit of measure called carbon dioxide equivalent (CO₂e) that allows for better comparison of the impact of different greenhouse gases. CO₂e is calculated by multiplying the mass amount of each greenhouse gas by its global warming potential (GWP) and then summing the resulting value. CO₂e was calculated using Equation 1:

Equation 1:

$$tons\ CO_2e = \sum_{i=0}^n GHG_i \times GWP_i$$

Where:

GHG_i = Mass emissions of each greenhouse gas

GWP_i = Global warming potential for each greenhouse gas

n = the number of greenhouse gases emitted

On November 29, 2013, the U.S EPA starting using the GWPs from the Intergovernmental Panel on Climate Change's (IPCC) Fourth Assessment Report (AR4) (IPCC 2007) in its programs and reports, including using it for the first time in the 2015 national greenhouse gas inventory for years 1990 – 2013.

In the past, DNR used the GWPs from Third Assessment Report (IPCC 2001) to calculate emissions in statewide Iowa Greenhouse Gas Inventories for years 2010 – 2015. However, starting with this report,

DNR used the IPCC AR4 GWPs for the first time and recalculated all emissions from 2005 – 2016. The GWP values used are shown in Table 2.

Table 2. Global Warming Potentials

Pollutant	GWP used by DNR (IPCC TAR 2001)	GWP used by EPA as of 11/29/13 (IPCC AR4 2007) and DNR as of 3/22/17
Carbon Dioxide (CO ₂)	1	1
Methane (CH ₄)	21	25
Nitrous Oxide (N ₂ O)	310	298
Sulfur Hexafluoride (S _F 6)	23,900	22,600
Hydrofluorocarbons (HFC)	Vary by pollutant – For a complete list, refer to DNR's Greenhouse Gas Emissions Estimation Guidance .	
Perfluorocarbons (PFC)		

Chapter 2 - Agriculture

This chapter includes non-energy greenhouse gas (GHG) emissions from livestock and crop production in Iowa. GHG emissions from fossil fuel-fired agricultural equipment are discussed in *Chapter 6 – Transportation* and carbon emissions and sinks from agriculture are discussed in *Chapter 9 – Land Use, Land Use Change, and Forestry (LULUCF)* of this document.

GHG emissions are emitted from four agricultural sectors in Iowa – enteric fermentation, manure management, agricultural soils, and agricultural burning. The GHGs emitted are methane (CH₄) and nitrous oxide (N₂O). Table 3 summarizes the source of GHG emissions in each sector. N₂O emissions from rice cultivation were not included, as rice is not grown in Iowa (USDA 2017b).

Table 3: Sources of Agricultural GHG Emissions in Iowa

Sector		GHGs Emitted	Source of Emissions
Enteric Fermentation		CH ₄	Microbial activity in the digestive systems of dairy cattle, beef cattle, sheep, goats, swine, and horses.
Manure Management		CH ₄ N ₂ O	Decomposition of manure during storage and treatment of livestock manure.
Agricultural Soils	Residues, legumes, and histosols	N ₂ O	Biological nitrogen fixation by crops, crop residues remaining on fields, and cultivation of high organic content soils (histosols).
	Fertilizers	N ₂ O	Application of manure, fertilizers, etc. to soils and leaching/runoff of nitrogen into ground or surface water.
	Animals	N ₂ O	Animal excretions directly on to soils such as pastures.
Agricultural Burning		CH ₄ N ₂ O	Burning of crop residues.

Method

GHG emissions from agriculture were calculated using the United States Environmental Protection Agency's (EPA) State Greenhouse Gas Inventory Tool (SIT) agriculture module dated January 3, 2017 (ICF 2017a and 2017b).

Enteric Fermentation

The SIT calculates CH₄ emissions from enteric fermentation by multiplying various livestock populations by an annual CH₄ emission factor (kilograms CH₄ per head). The data sources for the animal populations used are listed in Table 4. The number of "Feedlot Heifers" and "Feedlot Steers" was derived by applying a 35/65 heifer/steer ratio to the "Total Number on Feed".

Manure Management

This sector includes CH₄ and N₂O emissions from manure when it is being stored and treated in a manure management system. In general, CH₄ emissions increase in more anaerobic (lacking oxygen) conditions while N₂O emissions increase under aerobic conditions (Strait et al. 2008). The same dairy cattle, beef cattle, sheep, goat, swine, and horse populations were used as for the enteric fermentation sector for consistency. Several other animal types were added as shown in Table 4.

Table 4: Animal Populations

Animal Type	Year	Data Source
Dairy cattle	2016	2016 Iowa Agricultural Statistics Bulletin (USDA 2016)
Beef cattle		
Sheep		
Goats	2012 used as proxy for 2013 - 2016	USDA-NASS Quick Stats (USDA 2017b)
Horses		
Breeding swine	2016	2016 Iowa Agricultural Statistics Bulletin (USDA 2016)
Market swine under 60 lbs. ²		
Market swine 60 – 119 lbs. ³		
Market swine 120 – 179 lbs.		
Market swine over 180 lbs.	2012 used as proxy for 2013-2016	USDA-NASS Quick Stats (USDA 2017b)
Broilers		
Chickens		
Hens		
Pullets		
Turkeys		

In addition, the number of “Sheep on Feed” and “Sheep off Feed” were derived by applying a 6.5/93.5 on feed/off feed ratio to the total number of sheep.

Agricultural Residue Burning

Burning of cropland is not a typical agricultural practice in Iowa. According to Iowa State University Extension and Outreach,

“Burning corn and soybean fields is just NOT a practice that is used in Iowa or many other Midwest states as a way of preparing the fields for planting a subsequent crop. Yes, there are rare occasions where corn residue is burnt off a field but it would not even be 1% of the crop acres. An example would be if the residue washed and piled up in an area it may be burnt to allow tillage, planting and other practices to occur. Another rare occasion is when accidental field fires occur during harvesting of the corn crop. But again this would be less than 1% of the crop acres.” (Licht 2015).

The SIT over-estimates agricultural fires, as it assumes that 3% of Iowa corn, soybean, and wheat field residue is burned annually. The *Year 2000 Iowa Greenhouse Gas Emissions Inventory* notes that “According to expert opinion, even this lower estimate [3%] is thought to be too large in Iowa because

² SIT uses the category of market swine under 60 lbs., but USDA uses the category of market swine under 50 lbs.

³ SIT uses the category of market swine 60 – 119 lbs., but USDA uses the category of market swine 50 - 119 lbs.

burning is mostly a maintenance tool for conservation plantings, which are not extensive” (Wollin and Stigliani 2005). The DNR has been working with EPA emission inventory staff for several years to refine estimates for agricultural fires in the EPA’s National Emissions Inventory (NEI) and the DNR’s annual greenhouse gas inventories (DNR 2015, Pouliot 2015 and Stein 2015).

In 2015, DNR staff reviewed the details of 1,008 fires in 2014 that were reported to Iowa DNR by local fire departments (Kantak 2015) as shown in Table 5. Staff found that:

- 39 of the fires were truly agricultural fires, with 38 of 39 being fires being purposely set on grasslands enrolled in the Conservation Reserve Program, and one fire in a field of millet. No cornfield or soybean field fires were reported to DNR.
- 309 of the fires were identified as being prescribed fires (fires ignited by management actions to meet specific objectives): 166 on state land, 101 on private land, 37 on county land, 5 on federal land.
- 660 of the fires were identified as being wildfires. Seven were accidental fires in cornfields that were started by overheated harvesting equipment. Several were wildfires that spread when trash or brush burning spread out of control to a nearby field or ditch.

Table 5: Fires in 2014 Reported to Iowa DNR


Type of Fire	No. of Fires in 2014 Reported to Iowa DNR	Total Acres Reported	Average Acres Burned
Agricultural Fires	39	1,981.4	50.8
Prescribed Fires	309	14,701.7	47.6
Wildfires	660	12,218.6	18.5
Total	1,008	28,901.7	28.7

There are several discrepancies between the pollutants EPA calculates for agricultural fires in the NEI (EPA 2015) and the SIT (ICF 2016a). EPA calculates carbon dioxide (CO₂) and methane (CH₄) emissions in the NEI, but calculates emissions from methane (CH₄) and nitrous oxide (N₂O) in the SIT. In addition, the NEI calculates emissions from the burning of grasses and CRP lands, but the SIT only calculates emissions from crops. EPA calculates emissions from the burning of grass and pasture lands in the national GHG inventory, but not from crops (EPA 2016). Due to these discrepancies, emissions from agricultural residue burning were not included in this inventory. Resolving this discrepancy continues to be an area of future improvement in the inventory.

Agricultural Soils

N₂O emissions in the agricultural soils sector occur from many different pathways as shown in Figure 1 (EPA 2016). N₂O is emitted when the natural processes of denitrification and nitrification interact with agricultural practices that add or release nitrogen (N) in the soil profile. Denitrification is the process of converting nitrate to nitrogen gas. It is carried out by microorganisms in an oxygen-lacking environment. Nitrification occurs when ammonia is converted to nitrites and nitrates by naturally occurring, specialized bacteria in the environment.

Figure 1: Sources and Pathways of N₂O Emissions in Ag Soils (EPA 2016)



Direct N₂O emissions occur at the site of application of both synthetic and organic fertilizers to the soil, production of N-fixing crops, and integration of crop residues into the soil by practices such as cultivation. Indirect emissions occur when N is made available or is transported to another location following volatilization, leaching or runoff, and is then converted to N₂O (EPA 2016).

Plant Residues and Legumes

Crop production data for alfalfa, corn for grain, oats, soybeans, and wheat (USDA 2017b) was used to calculate N₂O from nitrogen-fixing crops, including alfalfa and soybeans, and nitrogen returned to soils during the production of corn for grain, wheat, oats, and soybeans.

Soil Cultivation - Nitrous Oxide (N₂O)

N₂O is also emitted during the cultivation of highly organic soils called histosols. May 2011 soil survey data from the Natural Resources and Conservation Service shows there are just over 70,000 acres of histosols in Iowa (Sucik 2011a and 2011b). The quantity of histosols that are cultivated is not currently available (Bedmarek 2012), so the DNR estimated the number of cultivated histosols acres by multiplying the acres of histosols by the annual percentages of Iowa cropland that are corn and soybeans (USDA 2017b) and by the average percentage of each crop that is tilled (Sucik 2011b). However, this may be an overestimation as according to former State Soil Scientist, Michael Sucik, "...all Histosols are listed as hydric soils and are eligible for the Wetland Restoration Program as CRP [Conservation Reserve Program] practices that require wetlands. Also, a Histosol would require some type of artificial drainage in order to be consistently row cropped" (Sucik 2011a).

Soil Tillage Practices

Carbon may be emitted when soils are tilled. However, carbon may also be sequestered when soil conservation practices are used (no-till or reduced tillage), are converted to the Conservation Reserve Program, or are converted grass, trees or wetlands. This balance between emissions and sequestration is called the soil carbon flux. The SIT does not include the ability to calculate emissions from soil carbon flux from tillage practices.

Practicing no-till for many consecutive years produces the greatest carbon sequestration. When soil is tilled, the soil becomes oxygenated, increasing microbial activity and releasing stored carbon. However, the amount of carbon stored and released is uncertain. Scientific studies and literature reviews, such as those by Baker et al. (2007), Blanco-Canqui, and Lal (2008), have created uncertainty in this area, while other studies such as those by Franzluebbers (2009) and Boddey et al (2009) dispute them. According to the USDA's "*No-Till Farming is a Growing Practice*", there is much uncertainty in the interaction between tillage practices, carbon, and other greenhouse gases" (USDA 2010). A 2007 study by West and Six explains that, "*The extent to which soil C accumulation occurs after a reduction in tillage intensity is determined by the history of land management, soil attributes, regional climate, and current carbon stocks*" (West and Six 2007). The relationship between tillage and nitrogen oxides (N₂O) is also not completely

certain. Several studies have observed increases, decreases, and no change in N₂O when soil is tilled (USDA 2010).

The complexity of calculating soil carbon flux is described in USDA's *Science-Based Methods for Entity-Scale Quantification of Greenhouse Gas Sources and Sinks from Agriculture and Forestry Practices*. This 605-page document was developed to create "a standard set of GHG estimation methods for use by USDA, landowners, and other stakeholders to assist them in evaluating the GHG impacts of their management decisions" (Eve 2014). It recommends that soil organic carbon stocks are calculated by modeling with the DAYCENT model. At this time, the DNR does not have the required data inputs or capability of running the DAYCENT model.


The USDA has also established seven regional climate change offices, offering climate hazard and adaptation data and services to farmers, ranchers, and forest landowners. The NRCS, a department within the USDA, has also launched a program called Carbon Management and Evaluation Online Tool (COMET-FARM) that allows users to calculate how much carbon is removed from the atmosphere from certain conservation efforts. The COMET-FARM website explains that:

The tool guides you through describing your farm and ranch management practices including alternative future management scenarios. Once complete, a report is generated comparing the carbon changes and greenhouse gas emissions between your current management practices and future scenarios (NRCS 2015).

COMET-FARM is not designed to calculate statewide greenhouse gas emissions from farming and ranching. It requires specific data inputs for each individual farm. However, if NRCS should publish results from the tool in the future, the DNR may include them in future inventory reports.

While the DNR is unable to quantify agricultural soil carbon flux at this time, it is known that cumulative Iowa acres in the CRP program are trending downward as shown in Figure 2. This indicates that the amount of carbon stored in agricultural soils *may* be decreasing as more soil is tilled each year. However, any effects from cover crops were not considered. This may be a future inventory improvement.

Figure 2: Acres Enrolled in CRP by Fiscal Year (USDA 2017a)



Fertilizer Utilization

The DNR calculated fertilizer emissions for 2016 using fertilizer tonnages from the Iowa Department of Agriculture and Land Stewardship's (IDALS) *Fertilizer Tonnage Distribution in Iowa* report (IDALS 2017). The IDALS fertilizer data is provided per the 2016 growing season, which is from July 2015 – June 2016. The 2016 growing season was then used as a proxy for the 2017 growing season (July 2016 – June 2017).

Adjustments

Emissions from 2005 - 2015 were recalculated using GWPs from IPCC AR4 as noted in the *General Calculation Method* section of this document. In addition, 2015 emissions from enteric fermentation, manure management, and agricultural soils have been updated since the DNR's 2015 GHG Inventory Report was published in December 2016 using revised activity data (such as animal populations or fertilizer application) from USDA or IDALS.


Results

GHG emissions from agriculture increased 1.82% from 2015 – 2016 and 17.51% from 2005 – 2016. Gross GHG emissions from agriculture were 39.48 MMtCO₂e in 2016, or 30.78% of Iowa's total gross GHG emissions. This total does not account for any carbon sinks from agriculture. Sinks are discussed in *Chapter 9 – Land Use, Land Use Change, and Forestry*. The majority of agricultural emissions (53.79%) are from soils as shown in Table 6 and Figure 3.

Table 6: Gross GHG Emissions from Agriculture (MMtCO_{2e})⁴

Category	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Enteric Fermentation	7.08	7.55	7.88	8.05	8.02	8.39	8.41	8.36	8.36	8.16	8.31	8.38
Manure Management	7.85	7.88	8.68	9.52	9.59	8.83	9.53	9.62	9.84	9.70	10.04	9.86
Agricultural Soils	18.66	20.28	23.67	19.07	18.87	19.16	19.98	18.69	18.82	19.90	20.43	21.24
Total	33.60	35.71	40.23	36.65	36.48	36.38	37.91	36.67	37.03	37.76	38.78	39.48

Figure 3: Gross GHG Emissions from Agriculture (MMtCO_{2e})



Enteric Fermentation

CH₄ emissions from enteric fermentation were 8.38 MMtCO_{2e} in 2016, increasing 0.83% from 2015. This can be attributed to a 1.97% increase in the total cattle population. The total swine population decreased 2.63%.


While poultry and swine make up the greatest percentages of total livestock in Iowa as shown in Figure 4, enteric fermentation emissions are primarily driven by the cattle population. This is because cattle emit more CH₄ than other ruminant animals due to their unique stomachs. In addition, poultry do not emit methane through enteric fermentation. The amount of methane emitted from each animal type is shown in Table 7.

⁴ Totals may not equal the sum of subtotals shown in this table due to independent rounding.

Table 7: Methane Emitted per Animal

Animal Type	kg/head CH ₄ Emitted (ICF 2016a)
Beef Cattle	42.0 – 95.1
Dairy Cattle	43.5 – 132.4
Goats	5.0
Horses	18.0
Sheep	8.0
Swine	1.5

Figure 4: 2016 Animal Populations (USDA 2016, 2017b)⁵



Manure Management

Factors influencing CH₄ and N₂O emissions include the animal type, animal population, animal mass, the type of manure management system, etc. GHG emissions from manure management decreased 1.74% from 2016 and accounted for 24.98% of agricultural GHG emissions in 2016. The decrease in emissions in 2015 can be linked to a decrease of 560,000 swine and 13,000 poultry produced in 2016 (USDA 2016 and USDA 2017b).


Agricultural Soils

The majority of GHG emissions from agricultural soils can be attributed to crop production (fertilizers, crop residues, and nitrogen fixing) as shown in Figure 5. N₂O emissions from agricultural soils increased 3.97% from the previous year due to:

⁵ The goat, horse, and sheep population each account for less than 1% of the total animal population.

- A 3.12% increase in nitrogen fertilizers applied to crop grounds.
- An 8.94% increase in field crop production (corn, oats, soybeans, and wheat).

Figure 5: 2015 - 2016 Gross GHG Emissions from Agricultural Soils (metric tons N₂O)



N₂O emissions from agricultural soils accounted for 53.79% of all agricultural GHG emissions and 16.56% of total statewide GHG emissions in 2016. The majority of GHG emissions from agricultural soils can be attributed to crop production (fertilizers, crop residues, and nitrogen fixing) as shown in Figure 5.

Table 8: Iowa Crop Production 2015 – 2016 (USDA 2017b)

Crop	2015 (1000 Bushels)	2016 (1000 Bushels)
Barley	85	85
Corn for Grain	2,505,600	2,740,500
Oats	4,161	3,268
Rye	46	46
Sorghum for Grain	59	59
Soybeans	533,700	571,725
Wheat	780	1,071
Total	3,044,430	3,319,063
Crop	2015 (1000 tons)	2016 (1000 tons)
Alfalfa	3,003	2,310

Uncertainty

Excerpted from SIT Agriculture Module (ICF 2017a):

Enteric Fermentation

The quantity of methane (CH₄) emitted from enteric fermentation from livestock is dependent on the estimates of animal populations and the emission factors used for each animal type. Therefore, the uncertainty associated with the emission estimate stems from those two variables. Uncertainty is also introduced as animal populations vary throughout the year. There is also uncertainty associated with the original population survey methods used by USDA. Emission factors vary in each animal, depending on its production and diet characteristics, as well as genetics (ICF 2017a).

Manure Management

As with enteric fermentation, uncertainty occurs in animal populations and the emission factors used for each animal. However, the largest contributor to uncertainty in manure management emissions in the SIT is the lack of lowa-specific data describing manure management systems and the CH₄ and N₂O emission factors used for these systems. In addition, there is uncertainty in the maximum CH₄ producing potential (B₀) used for each animal group. This value varies with both animal and diet characteristics, so estimating an average across an entire population introduces uncertainty. While the B₀ values used in the SIT vary by animal subcategory to attempt to represent as many of these differences as possible, there is not sufficient data available at this time to estimate precise values that accurately portray the B₀ for all animal types and feeding circumstances (ICF 2004).

Agricultural Soils

The amount of N₂O emission from managed soils is dependent on a large number of variables other than N inputs. They include soil moisture, pH, soil temperature, organic carbon availability, oxygen partial pressure, and soil amendment practices. The effect of the combined interaction of these variables on N₂O flux is complex and highly uncertain. The methodology used in the SIT is based only on N inputs, does not include other variables, and treats all soils, except histosols, equally. In addition, there is limited knowledge regarding N₂O productions from soils when N is added to soils. It is not possible to develop emission factors for all possible combinations of soil, climate, and management conditions.

Uncertainties also exist in fertilizer usage calculations. The fertilizer usage does not include non-commercial fertilizers other than manure and crop residues, and site-specific conditions are not considered in determining the amount of N excreted from animals. Additional uncertainty occurs due to lack of lowa-specific data for application of sewage sludge and cultivation of histosols (ICF 2017a).

Uncertainties in the estimation method for agricultural residue burning are noted above under the "Methods" heading.

Chapter 3 – Fossil Fuel Consumption

This chapter includes GHG emissions from fossil fuel consumption in four categories: power plants, residential, industrial, and commercial. The residential, industrial, and commercial categories are often combined into one category called RCI. Fossil fuels combusted by mobile sources are included in the transportation sector and discussed later in this report in *Chapter 6 – Transportation*. Emissions from the electric generation category include direct emissions resulting from the combustion of fossil fuels at the electric generating station (i.e. power plant). Indirect emissions from electricity consumed at the point of use (i.e. residential electric water heaters) are discussed in *Chapter 10 – Indirect Emissions from Electricity Consumption*.

Method

Residential, Commercial, Industrial (RCI)

GHG emissions were calculated using two SIT modules – the CO₂FFC module for carbon dioxide (CO₂) emissions and the Stationary Combustion module for CH₄ and N₂O emissions (ICF 2017a-d). These modules calculate energy emissions based on annual statewide consumption for the sectors and fuels listed in Table 9:

Table 9: Fuel Types Included in Fossil Fuel Consumption

Fuel Types	Residential	Commercial	Industrial
Asphalt/Road oil			X
Aviation gasoline blending components			X
Coal	X	X	X
Coking coal, other coal			X
Crude oil			X
Distillate fuel oil	X	X	X
Feedstocks			X
Kerosene	X	X	X
LPG	X	X	X
Lubricants			X
Misc. petroleum products			X
Motor gasoline		X	X
Motor gasoline blending components			X
Natural gas	X	X	X
Pentanes plus			X
Petroleum coke			X
Residual fuel		X	X
Still gas			X
Special naphthas			X
Unfinished oils			X
Waxes			X
Wood	X	X	X

Iowa-specific 2016 energy consumption data will not be published by the U.S. Energy Information Administration until June 2018, so the DNR projected 2016 energy consumption. This was done by using the EIA's *Annual Energy Outlook (AEO) 2017 with Projections to 2050* (EIA 2017a) and 2015 bulk energy consumption data from the EIA's State Energy Data System (SEDS) (EIA 2017b). The AEO2017 includes several different projection cases, which each address different uncertainties. The DNR used the AEO2017 "Reference Case", which models projections of what may happen given certain assumptions and methodologies. The AEO uses the National Energy Modeling System (NEMS), which has the objective to show various interactions of economic changes and energy supply, demand, and prices (EIA 2017a). The Reference Case assumes implementation of the Clean Power Plan (CPP). However, the CPP is currently stayed, and on October 10, 2017, the EPA proposed to repeal the CPP. The uncertainty with the regulation adds a level of uncertainty to the DNR's projections.

The projections in the Reference Case are done at the regional level, and Iowa is in the West North Central U.S. Census Region. The 2016 energy consumption was estimated for each fuel type using one of two methods as described below and shown in Table 10:

Fuel Method 1

The percent change in the regional consumption of each fuel type in the AEO2017 was calculated. The percent change was then applied to the Iowa 2015 fuel consumption in SEDS. This method was used for the fuel types listed in Table 10. This method is different from previous years where the ratio of Iowa fuel consumption from SEDS to the regional fuel consumption for the previous year in the AEO was calculated and then applied to the predicted regional fuel consumption for the current year in the AEO.

Fuel Method 2

These sectors were not included in the AEO Reference Case, so it was assumed that 2016 fuel consumption was equal to the 2015 fuel consumption. This method was used for the fuel types listed in Table 10.

Table 10: Method Used to Estimate 2016 Fuel Consumption

Fuel Type	Estimation Method
Commercial Distillate Fuel Oil	Method 1
Commercial Kerosene	Method 1
Commercial LPG	Method 1
Commercial Motor Gasoline	Method 1
Commercial Natural Gas	Method 1
Industrial Coal	Method 1
Industrial Distillate Fuel Oil	Method 1
Industrial LPG	Method 1
Industrial Motor Gasoline	Method 1
Industrial Natural Gas	Method 1
Industrial Other Coal	Method 1
Residential Distillate Fuel	Method 1
Residential LPG	Method 1
Residential Natural Gas	Method 1

Table 10 (continued)

Fuel Type	Estimation Method
Industrial Crude Oil	Method 2
Commercial Coal	Method 1
Commercial Residual Fuel	Method 2
Commercial Wood	Method 2
Industrial Asphalt and Road Oil	Method 2
Industrial Aviation Gasoline Blending Components	Method 2
Industrial Coking Coal	Method 2
Industrial Feedstocks, Naphtha less than 401 F	Method 2
Industrial Feedstocks, Other Oils greater than 401 F	Method 2
Industrial Kerosene	Method 2
Industrial Lubricants	Method 2
Industrial Misc. Petro Products	Method 2
Industrial Motor Gasoline Blending Components	Method 2
Industrial Pentanes Plus	Method 2
Industrial Petroleum Coke	Method 2
Industrial Residual Fuel	Method 2
Industrial Special Naphthas	Method 2
Industrial Still Gas	Method 2
Industrial Unfinished Oils	Method 2
Industrial Waxes	Method 2
Industrial Wood	Method 2
Residential Coal	Method 2
Residential Kerosene	Method 2
Residential Wood	Method 2

Power Plants

Emissions from emissions from electricity generation at power plants were not calculated using fuel consumption data. Depending on the year, emissions from either EPA's federal Acid Rain program (CAMD 2016) or federal GHG Reporting Program (40 CFR 98, EPA 2016) were used as follows:

2005 – 2009

CO₂ emissions reported to EPA by individual facilities subject to the Acid Rain program were used. This data is more accurate than the values from EIA because the CO₂ emissions reported by facilities to EPA are actual measured emissions values from continuous emission monitors (CEMS) located on electric generating units.

2010 - 2016

Power plants became subject to the federal GHG reporting program starting with calendar year 2010. Facilities are required to report CO₂, CH₄, and N₂O emissions. Like the Acid Rain program, this CO₂ data is from CEMS and is more accurate than EIA data. In addition, the CH₄ and N₂O emissions are calculated using facility-specific fuel heating values. The CO₂ data reported to the federal GHG reporting program

is consistent with the CO₂ emissions reported by the same facilities to EPA as required by the Acid Rain Program.

Adjustments

As noted in the *General Calculation Method* section of this document, emissions from 2005 - 2015 were recalculated using GWPs from IPCC AR4. In addition, the DNR previously forecasted 2015 emissions from RCI due to a lack of low-specific energy consumption data. However, the 2015 energy data was released by EIA in June 2017 (EIA 2017b), so the DNR used the data to recalculate 2015 emissions as shown in Table 11.

Table 11: Recalculated RCI Emissions (MMtCO₂e)

Category	2015 Value Published Dec. 2016	2015 Updated Value
Residential	4.28	4.49
Commercial	4.42	4.60
Industrial	22.25	22.44
Total	30.95	31.54

Results

Total GHG emissions from fossil fuel consumption in 2016 were 57.23 MMtCO₂e, a decrease of 3.77% from 2015 and 10.39% from 2005 levels as shown in Table 12 and Figure 6. Emissions from each of the three RCI fossil fuel categories increased in 2016 due to increased fossil fuel consumption:

- residential fuel use emissions increased 1.27%
- commercial fuel use emissions increased 3.52%
- industrial fuel use emissions increased 0.65%


Alternately, emissions from power plants (electric generating facilities) decreased 14.02% from 2015 and have decreased 31.23% from 2005. This is because the amount of electricity generated from zero-emitting sources such as wind and solar and low-emitting sources such of natural gas is increasing.

Table 12: GHG Emissions from Fossil Fuel Consumption by Category (MMtCO₂e)⁶

Category/ Fuel Type	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Residential	4.89	4.55	4.89	5.58	5.21	4.88	4.85	4.01	5.12	5.38	4.49	4.55
Commercial	3.49	3.85	3.95	4.42	4.70	4.48	4.52	4.11	4.83	4.92	4.60	4.76
Industrial	18.65	18.86	19.72	21.34	20.82	21.88	22.07	21.84	22.87	22.52	22.44	22.59
Power Plants	36.83	36.35	40.04	41.78	37.71	42.33	38.98	35.76	33.06	33.44	29.46	25.33
Total	63.87	63.61	68.59	73.12	68.44	73.56	70.42	65.72	65.89	66.26	61.00	57.23

⁶ Values do not include emissions from the transportation sector. Totals may not equal the sum of subtotals shown in this table due to independent rounding.

Figure 6: GHG Emissions from Fossil Fuel Consumption by Category (MMtCO₂e)



Uncertainty -

CO₂ Emissions - Excerpted from SIT CO₂FFC Module (ICF 2017a):

The amount of CO₂ emitted from energy consumption depends on the type and amount of fuel that is consumed, the carbon content of the fuel, and the fraction of the fuel that is oxidized. Therefore, the more accurate these parameters are, the more accurate the estimate of direct CO₂ emissions will be. Nevertheless, there are uncertainties associated with each of these parameters.

National total energy consumption data is fairly accurate, but there is more uncertainty in the state-level data, especially when allocating consumption to the individual end-use sectors (i.e. residential, commercial, and industrial). The amount or rate at which carbon is emitted to the atmosphere can vary greatly depending on the fuel and use, and may vary at the state-level compared to the national default levels in the SIT.

The uncertainty in carbon content and oxidation are much lower than with fuel consumption data. Carbon contents of each fuel type are determined by EIA by sampling and the assessment of market requirements, and, with the exception of coal, do not vary significantly from state to state. EIA takes into account the variability of carbon contents of coal by state; these coefficients are also provided in the SIT.

Uncertainty is also introduced by the complexity in calculating emissions from the import/export of electricity. The precise fuel mix used to generate the power crossing state lines is very difficult to determine, so, an average fuel mix for all electricity generation within a specific region of the grid must usually be used. Moreover, these emissions factors are generated by emission monitors (rather than carbon contents of fuels), which may overestimate CO₂ emissions to a small extent (ICF 2017a).

CH₄ and N₂O Emissions - Excerpted from SIT Stationary Combustion Module (ICF 2017b):

The amount of CH₄ and N₂O emitted depends on the amount and type of fuel used, the type of technology in which it is combusted (e.g., boilers, water heaters, furnaces), and the type of emission control used. In general, the more detailed information available on the combustion activity, the lower the uncertainty. However, as noted in the Revised 1996 IPCC Guidelines (IPCC/UNEP/OECD/IEA 1997), the contribution of CH₄ and N₂O to overall emissions is small and the estimates are highly uncertain.

Uncertainties also exist in both the emission factors and the EIA energy consumption data used to calculate emissions. For example, the EIA state data sets do not fully capture the wood used in fireplaces, wood stoves, and campfires. As with CO₂, uncertainty is also introduced with allocating energy consumption data to the individual end-use sectors and estimation of the fraction of fuels used for non-energy (ICF 2017b).

Chapter 4 - Industrial Processes

This chapter includes non-combustion GHG emissions from a variety of industrial processes. The processes and GHG pollutants emitted from each category are shown in Table 13. Emissions from these industries do not include emissions from fossil fuel combustion, which are included in *Chapter 3 – Fossil Fuel Combustion*.

Table 13: Industrial Processes and GHG Emissions

Category	GHGs Emitted
Cement Production	CO ₂
Lime Manufacture	CO ₂
Limestone and Dolomite Use	CO ₂
Soda Ash Use	CO ₂
Iron and Steel Production	CO ₂
Ammonia Production & Urea Consumption	CO ₂
Nitric Acid Production	N ₂ O
Ozone Depleting Substances (ODS) Substitutes	HFCs, PFCs, and SF ₆
Electric Power Transmission and Distribution	SF ₆

Cement Production

Carbon Dioxide (CO₂) is emitted during a process called calcining when limestone is heated in a cement kiln to form lime and CO₂. The CO₂ is vented to the atmosphere and the lime is then mixed with silica-containing materials such as clay to form clinker, an intermediate product that is made into finished Portland cement (ICF 2004). Two facilities in Iowa currently produce Portland cement.

Lime Manufacture

Similar to cement manufacturing, lime is produced by heating limestone in a kiln, creating lime and CO₂. The CO₂ is typically released to the atmosphere, leaving behind a product known as quicklime, which can then be used to produce other types of lime (ICF 2004). One facility currently manufactures lime in Iowa.

Limestone and Dolomite Use

Limestone and dolomite are used in industrial processes such as glass making, flue gas desulfurization, acid neutralization, etc.

Soda Ash Use

Soda ash is currently only produced in three states – Wyoming, Colorado, and California. However, commercial soda ash is used as a raw material in a variety of industrial processes and in many familiar consumer products such as glass, soap, and detergents (ICF 2017b). In Iowa, it is commonly used by corn wet millers for pH control, in ion exchange regeneration, and in other operations (DNR 2010).

Iron and Steel

Iron and steel production is an energy-intensive process that also generates process-related GHG emissions. Steel is produced from pig iron or scrap steel in a variety of specialized steel-making furnaces, including electric arc furnaces (EAFs) and basic oxygen furnaces (BOFs) (EPA 2016). There are currently no pig iron mills operating in Iowa. All three steel production facilities currently operating in Iowa use EAFs to produce steel from scrap. These furnaces use carbon electrodes, coal, natural gas, and other substances such as limestone and dolomite to aid in melting scrap and other metals, which are then improved to create the preferred grade of steel. In EAFs, CO₂ emissions result primarily from the consumption of carbon electrodes and from the consumption of supplemental materials used to augment the melting process (EPA 2016).

Ammonia Production and Urea Consumption

CO₂ is released during the manufacture of ammonia. The chemical equations to calculate the release of CO₂ are fairly complicated, but in general, anhydrous ammonia is synthesized by reacting nitrogen with hydrogen. The hydrogen is typically acquired from natural gas. The majority of direct CO₂ emissions occur when the carbon in the natural gas is then eliminated from the process by converting it to CO₂. Other emissions of CO₂ can occur during condensate stripping or regeneration of the scrubbing solution. CO₂ emissions may also be captured for use in urea synthesis or carbon sequestration and storage (WRI 2008). Three facilities in Iowa currently produce ammonia.

Nitric Acid Production

Nitrous Oxide (N₂O) is produced when ammonia is oxidized to produce nitric acid. Two facilities in Iowa currently produce nitric acid.

Consumption of ODS Substitutes

Ozone Depleting Substances (ODS) are often used in refrigeration, air conditioning, aerosols, solvent cleaning, fire extinguishers, etc. However, ODS are being phased out per the Montreal Protocol and the 1990 Clean Air Act Amendments. The most common ODS are HFCs, but PFCs and SF₆ may also be used (ICF 2017b).

Electric Power Transmission and Distribution

SF₆ is used as an insulator in electricity transmission and distribution in equipment such as transformers, high-voltage circuit breakers, substations, and transmission lines (ICF 2017b).

Other Industry Types

GHG emissions from soda ash manufacturing, adipic acid production, (primary) aluminum production, HCFC-22 production, semiconductor manufacturing, and magnesium production and processing were not calculated as the DNR is not aware of any of these facilities currently operating in Iowa.

Method

2016 emissions from industrial processes were calculated using either the SIT (ICF 2017a) or using GHG emissions reported to EPA by individual facilities to the federal GHG reporting program (40 CFR 98, EPA 2017a) as shown in Table 14.

Table 14: Industrial Processes Calculation Methods and Activity Data

Category	Year	Calculation Method	Data Source
Ammonia and Urea Production	2016	40 CFR 98 Subpart G	(EPA 2017a)
Cement Production		40 CFR 98 Subpart H	(EPA 2017a)
Iron and Steel Production		40 CFR 98 Subpart Q	(EPA 2017a)
Lime Manufacture		40 CFR 98 Subpart S	(EPA 2017a)
Nitric Acid Production		40 CFR 98 Subpart V	(EPA 2017a)
Electric Power Transmission and Distribution	2015 as proxy for 2016	SIT	(EIA 2017), (EPA 2017b)
Limestone and Dolomite Use	2014 as proxy for 2015 and 2016	SIT	(USGS 2016)
ODS Substitutes		SIT	SIT default value
Soda Ash Use	2016	SIT	(USGS 2017)

Categories Calculated using the SIT

Emissions from use of limestone and dolomite in industrial processes were calculated by multiplying Iowa’s annual consumption by the ratio of national consumption for industrial uses to total national consumption.

Emissions from ODS substitutes and soda ash consumption categories were calculated by assuming that Iowa emissions were 0.97% of national emissions because Iowa’s population is 0.97% of the total U.S. Population (U.S. Census 2017).

Emissions from electric power transmission distribution were calculated by determining the ratio between 2015 Iowa retail sales vs. 2015 national retail sales (EIA 2017), and applying that ratio to 2015 national emissions of sulfur hexafluoride (SF₆). 2015 emissions were used as a proxy for 2016.

Adjustments

As noted in the *General Calculation Method* section of this report, emissions from Nitric Acid Production, Electric Power T & D, and ODS Substitutes were recalculated using GWPs from IPCC AR4.

Emissions from ODS Substitutes in 2015 were also recalculated using 2015 national emissions (EPA 2017b), adjusted for Iowa population (U.S. Census 2017). In the previous inventory, 2014 data was used as a proxy for 2015.

Table 15: Recalculated Emissions from Industrial Processes (MMtCO₂e)

Sector	2015 Value Published Dec. 2016	2015 Updated Value
ODS Substitutes	1.34	1.64


Results

GHG emissions from industrial processes in 2016 were 5.50MMtCO₂e, or 4.29% of total statewide GHG emissions. Emissions from this sector increased 4.28% from 2015 as shown in Table 16. Ammonia and urea production, ODS substitutes, nitric acid production, and cement manufacture were the highest contributors to industrial process emissions in 2015 as shown in Figure 7.

Table 16: GHG Emissions from Industrial Processes (MMtCO₂e)⁷

Category	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Ammonia & Urea	1.01	0.91	0.95	0.87	0.60	0.84	0.75	0.85	0.88	0.86	0.81	0.92
Cement Manufacture	1.27	1.29	1.27	1.31	0.84	0.72	0.79	1.27	1.41	1.38	1.50	1.58
Electric Power T&D	0.12	0.11	0.10	0.10	0.09	0.08	0.06	0.06	0.05	0.05	0.05	0.05
Iron & Steel Production	0.13	0.13	0.13	0.12	0.09	0.23	0.20	0.23	0.19	0.18	0.16	0.19
Lime Manufacture	0.18	0.17	0.16	0.17	0.13	0.18	0.18	0.18	0.16	0.17	0.13	0.15
Limestone & Dolomite Use	0.18	0.29	0.24	0.25	0.29	0.39	0.16	0.15	0.33	0.21	0.21	0.21
Nitric Acid Production	0.65	0.72	0.78	0.86	0.87	0.95	0.90	0.96	0.80	0.82	0.74	0.75
ODS Substitutes	1.00	1.06	1.13	1.21	1.30	1.39	1.43	1.47	1.51	1.57	1.64	1.63
Soda Ash Consumption	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Total	4.57	4.71	4.78	4.92	4.22	4.79	4.49	5.17	5.34	5.28	5.28	5.50

Figure 7: 2016 GHG Emissions from Industrial Processes (MMtCO₂e)



⁷ Totals may not equal the sum of subtotals shown in this table due to independent rounding.

Uncertainty

Uncertainty occurs in categories where SIT default activity data was used instead of lowa-specific activity data, such as limestone and dolomite use, soda ash use, ODS substitutes, and electric power transmission and distribution.

Other major sources of uncertainty associated with calculating emissions from industrial processes are listed below (*Excerpted from SIT Industrial Processes Module (ICF 2017a)*).

- The estimation of emissions for limestone and dolomite use contains some inherent uncertainty based on limestone's variable composition.
- The use of population to disaggregate national emissions adds significant uncertainty.
- Uncertainties in emission estimates for electric power transmissions and distribution can be attributed to apportioning national emissions based on electricity sales. This method incorporates a low probability assumption that various emission reduction practices by Industry occur evenly throughout the country.

Chapter 5 - Natural Gas Transmission & Distribution

This chapter includes GHG emissions from natural gas transmission and distribution (T & D) in Iowa. In this sector, methane (CH₄) is emitted from leaks, vents, regulators, valves, compressors, accidents, and other devices located along the natural gas transmission and distribution networks. Carbon dioxide (CO₂) emissions from venting and flaring was not calculated due to lack of data. GHG emissions from coal mining, natural gas production, oil production, oil transmission, and oil transportation are not included as those industries are currently not active in Iowa.

Method

Natural Gas Transmission

Natural gas is transmitted in Iowa through large, high-pressure lines. These lines transport natural gas from production fields and processing plants located out-of-state to Iowa storage facilities, then to local distribution companies (LDCs) and high volume customers. Compressor stations, metering stations, and maintenance facilities are located along the transmission system. CH₄ is emitted from leaks, compressors, vents, and pneumatic devices (ICF 2017b).

The number of miles of transmission pipeline in Iowa was obtained from the United States Department of Transportation (DOT) Pipeline and Hazardous Materials Safety Administration's (PHMSA) Office of Pipeline Safety (DOT 2017). The Iowa Utilities Board confirmed that the number of natural gas compressor and gas storage stations did not change from the previous year (Munyon 2017).

Natural Gas Distribution

Natural gas is distributed through large networks of small, low-pressure pipelines. Natural gas flows from the transmission system to the distribution network at municipal gate stations, where the pressure is reduced for distribution within municipalities. CH₄ is emitted from leaks, meters, regulators, and accidents (ICF 2017b). Activity data from the DOT PHMSA's Office of Pipeline Safety was used for calculating emissions (DOT 2017). Data entered included miles of steel and cast iron distribution pipeline, unprotected and protected; number of services; and number of steel services, unprotected and protected.

Natural Gas Venting and Flaring

The DNR is unable to find data on the annual amount of natural gas vented and flared from natural gas transmission pipelines. This data is not tracked by the EIA (Little 2011), and the DNR has previously requested, but not received, this information from the Federal Energy Regulatory Agency (FERC). Therefore, no GHG emissions were calculated from natural gas venting and flaring.

Adjustments

As noted in the *General Calculation Method* section of this document, emissions from 2005 - 2015 were recalculated using GWPs from IPCC AR4.


Results

Total GHG emissions from natural gas transmission and distribution were 1.4073 MMtCO₂e⁸ in 2016, an increase of 0.62% from 2015 and an increase of 2.86% from 2005 as shown in Table 17 and Figure 8. Emissions increased in 2016 due to an increase in the number of services (e.g. gas meters) in the state. GHG emissions from this sector account for 1.10% of 2016 statewide GHG emissions.

Table 17: GHG Emissions from Natural Gas T & D (MMtCO₂e)

Category	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Transmission	0.7707	0.7723	0.7844	0.7857	0.7868	0.7871	0.7858	0.7862	0.7865	0.7864	0.7868	0.7867
Distribution	0.5974	0.5983	0.6008	0.6095	0.6084	0.6031	0.6132	0.6158	0.6135	0.6168	0.6118	0.6205
Total	1.3681	1.3706	1.3852	1.3953	1.3952	1.3901	1.3990	1.4020	1.4000	1.4031	1.3986	1.4073

Figure 8: GHG Emissions from Natural Gas T & D (MMtCO₂e)



⁸ DNR uses two decimal places throughout this report for consistency. However, in this sector four decimal places are needed show the difference in emissions from year to year.

Uncertainty

Excerpted from SIT Natural Gas and Oil Systems Module (ICF 2017a):

The main source of uncertainty in the SIT calculation methods is the emission factors. The emission factors used are based on a combination of statistical reporting, equipment design data, engineering calculations and studies, surveys of affected facilities and measurements. In the process of combining these individual components, the uncertainty of each individual component is pooled to generate a larger uncertainty for the overall emission factor. In addition, statistical uncertainties arise from natural variation in measurements, equipment types, operational variability, and survey and statistical methodologies. The method also does not account for regional differences in natural gas infrastructure and activity levels (ICF 2017a).

Chapter 6 - Transportation

This chapter includes GHG emissions from both highway and non-highway vehicles such as aviation, boats, locomotives, tractors, other utility vehicles, and alternative fuel vehicles.

Method

Emissions were calculated using the SIT Mobile Combustion module (ICF 2017a), which was updated by EPA to in 2016 to calculate CO₂, CH₄, and N₂O emissions from highway vehicles based on vehicle miles traveled. This is a more accurate than previous methods as it accounts for the vehicle type and vehicle age in the calculation, as well as accounting for the annual vehicle miles traveled. Emissions from non-highway vehicles were calculated based on fossil fuel consumption.

Highway Vehicles (CH₄ and N₂O)

Highway vehicles include passenger cars, truck, motorcycles, and heavy-duty vehicles. CH₄ and N₂O emissions from highway vehicles were calculated using the SIT as follows:

1. The vehicle miles traveled (VMT) for each vehicle type was calculated using the total 2016 annual VMT of 33,263 million miles (IDOT 2017). Neither the IDOT nor FHWA track state-level VMT by the seven classes used in the SIT, so the VMT was distributed among seven vehicle/fuel classes using the national distribution percentages from the Tables A-96 and A-97 from Annex 3 of the most recent national GHG inventory, *Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-2015* (EPA 2017). The classes and the national distribution percentages are shown in Table 18.

Table 18: VMT Vehicle/Fuel Classes and Distribution

Vehicle Class	Acronym	2015 (EPA 2017)	2016 Iowa VMT (10 ⁶ miles)
Heavy duty diesel vehicle	HDDV	8.29%	2,759
Heavy duty gas vehicle	HDGV	1.03%	344
Light duty diesel truck	LDDT	0.77%	255
Light duty diesel vehicle	LDDV	0.33%	111
Light duty gasoline truck	LDGT	19.68%	6,546
Light duty gasoline vehicle	LDGV	69.26%	23,037
Motorcycle	MC	0.64%	212
Total		100.00%	33,263

2. The VMT was then converted for use with existing emission factors. Iowa-specific emission factors were not available, so the SIT default emission factors were used. These factors are consistent with those used in the most recent national GHG inventory.
3. Next, the VMT was allocated by model year. Iowa-specific VMT data by model year was not available, so the VMT was allocated using the default national on-road age distribution by vehicle/fuel type in the SIT. The "Annual Vehicle Mileage Accumulation" table in SIT was updated to match that in Table A-101 in the most recent national inventory (EPA 2017).

- The control technology was then allocated by model year. Iowa-specific control technologies by model year were not available, so the national control technology values were used. The values in the SIT matched the Tables A-104, A-105, and A-106 in Annex 3 of the most recent national inventory (EPA 2017). One hundred percent was used for Tier 2 vehicles for 2015 and 2016.

Non-highway Vehicles (CH₄ and N₂O)

Non-highway vehicles include aviation, marine vessels, locomotives, and tractors. In general, CH₄ and N₂O emissions from non-highway vehicles were calculated using data from either the Energy Information Administration (EIA) or Federal Highway Administration as shown in Table 19.

Table 19: Iowa-specific Non-Highway Activity Data Used

Vehicle Type	Fuel Type	Year	Data Source
Aviation	Gasoline	2015 used as proxy for 2016	EIA SEDS (EIA 2017b)
Aviation	Jet Fuel, Kerosene		
Boats	Gasoline	2015 used as proxy for 2016	FHWA 2017
Heavy Duty Utility			
Tractors			
Construction			
Locomotives	Distillate Fuel	2015 used as proxy for 2016	EIA Adjusted Sales (EIA 2017a)
Tractors			
Construction	Distillate Fuel	2014 used as proxy for 2015-2016	SIT default value
Heavy Duty	Distillate Fuel		
Small Utility	Gasoline		
Alternative Fuel Vehicles			

Adjustments

As noted in the *General Calculation Method* section of this document, emissions from 2005 - 2015 were recalculated using GWPs from IPCC AR4.

CO₂ emissions for 2005 - 2010 emissions were recalculated using the SIT Mobile Combustion module as shown in Table 20. In previous years, CO₂ emissions for 2005 – 2010 emissions were calculated using the SIT CO₂FFC module. The ability to calculate CO₂ emissions was added to the SIT Mobile Combustion module by EPA in 2016.

Table 20: Recalculated CO₂ Emissions from Transportation (MMtCO₂e)

Pollutant	Year	Value Published Dec. 2016	Updated Value
CO ₂	2005	21.26	19.50
	2006	21.82	19.78
	2007	22.32	19.46
	2008	21.55	19.54
	2009	21.13	19.00
	2010	21.56	19.04

Results

Total GHG emissions from transportation were 19.36 MMtCO₂e in 2016 as shown in Table 21. This is an increase of 0.28% from 2015. CO₂ is the most prevalent GHG, accounting for 99.02% of GHG emissions from the transportation sector.

Table 21: GHG Emissions from Transportation (MMtCO₂e)⁹

Pollutant	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
CO ₂	19.50	19.78	19.46	19.54	19.00	19.04	19.27	19.31	19.21	19.32	19.10	19.17
CH ₄	0.05	0.05	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.03
N ₂ O	0.59	0.52	0.46	0.40	0.36	0.33	0.27	0.24	0.21	0.19	0.17	0.16
Total	20.14	20.35	19.96	19.99	19.40	19.41	19.58	19.59	19.46	19.55	19.31	19.36

The majority of emissions (58.95%) are from gasoline highway vehicles as shown in Figure 9. The SIT shows that while CO₂ emissions vary from year to year, emissions of CH₄ and N₂O have steadily decreased as shown in Figure 10 and Table 22. Nationally, CH₄ emissions declined by 64% and N₂O emissions decreased 63.3% from 1990 - 2015, due to national emission limits and control technologies in on-road vehicles (EPA 2017).

Table 22: Total CH₄ and N₂O Emissions from Mobile Sources (MMtCO₂e)^{10, 11}

Fuel /Vehicle Type	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Gasoline Highway	0.5824	0.5152	0.4491	0.3858	0.3495	0.3095	0.2522	0.2219	0.1942	0.1761	0.1605	0.1436
Diesel Highway	0.0055	0.0061	0.0062	0.0062	0.0053	0.0048	0.0056	0.0047	0.0042	0.0038	0.0034	0.0029
Non-Highway	0.0468	0.0470	0.0447	0.0503	0.0462	0.0489	0.0462	0.0486	0.0467	0.0458	0.0403	0.0403
Alternative Fuels	0.0025	0.0024	0.0029	0.0028	0.0029	0.0034	0.0046	0.0049	0.0049	0.0049	0.0049	0.0049
Total	0.6372	0.5707	0.5029	0.4452	0.4039	0.3665	0.3086	0.2801	0.2500	0.2306	0.2092	0.1917

⁹ Totals may not equal exact sum of subtotals shown in this table due to independent rounding.

¹⁰ Ibid.

¹¹ DNR uses two decimal places throughout this report for consistency. However, in this sector additional decimal places are needed show the difference in CH₄ emissions from year to year.

Figure 9. 2016 GHG Emissions per Fuel/Vehicle Type





Figure 10: CH₄ and N₂O Emissions by Fuel and Vehicle Type (MMtCO₂e)



Uncertainty

Uncertainty occurs because national vehicle/fuel type, age distributions, and emission factors, which may not be reflective of Iowa conditions, were applied to Iowa-specific VMT data. There is also some uncertainty in the method EPA used to develop the national vehicle/fuel type distributions and to develop emission factors (EPA 2017). The VMT used for alternative fuel vehicles has a higher level of uncertainty because the DNR was unable to locate Iowa-specific VMT data. Uncertainty may be introduced if the fuel consumption data or emission factors used do not reflect Iowa scenarios, such as using default national emission factors. In addition, it is assumed that all fuel purchased is consumed in the same year (ICF 2017b).

Chapter 7 – Waste: Solid Waste

This chapter includes methane (CH₄) emissions from municipal solid waste landfills and carbon dioxide (CO₂) and nitrous oxide (N₂O) emitted from the combustion of municipal solid waste to produce electricity. CH₄ emissions from landfills are a function of several factors, including:

- The total quantity of waste in municipal solid waste landfills,
- The characteristics of the landfills such as composition of the waste, size, climate; the quantity of CH₄ that is recovered and either flared,
- The quantity of CH₄ oxidized in landfills instead of being released into the atmosphere.

Fluctuations in CH₄ emissions can be caused by changes in waste composition, the quantity of landfill gas collected and combusted, the frequency of composting, and the rate of recovery of degradable materials such as paper and paperboard (EPA 2011).

Method

Municipal Solid Waste (MSW) Landfills

The DNR used emissions reported by MSW landfills to the EPA GHGRP (EPA 2017), which are calculated based on the characteristics of each individual report. EPA requires MSW landfills that emit 25,000 metric tons CO₂e or more to report their emissions. This included twenty-two Iowa landfills in 2016. An additional twenty-five Iowa MSW landfills were not required to report to the GHGRP. To calculate emissions for those that did not report to the GHGRP, the DNR calculated the potential methane emissions using EPA's Landfill Gas Emissions Model (LandGEM) version 3.02. It is based on a first-order decomposition rate equation for quantifying emissions from the decomposition of landfilled waste in MSW landfills (EPA 2005).

Combustion of Municipal Solid Waste

The amount of CH₄ emitted from power plants burning MSW to produce electricity was calculated using data reported annually by individual facilities to the DNR's Air Quality Bureau on their annual air emissions inventories. One facility reported burning a total of 23,950 tons of municipal solid waste in 2016 (Trower 2017). The DNR used state-specific proportions of discards that are plastics, synthetic rubber, and synthetic fibers instead of SIT default values to calculate CO₂ emissions from MSW combustion. These state-specific proportion values are from the 2011 Iowa Statewide Waste Characterization Study (MSW 2011). The state-specific proportions of discards used are shown in Table 23.

Table 23: Proportions of Discards used in the Solid Waste Module

Material	SIT Default Value	2011 Iowa Study
Plastics	17.0 – 18.0%	16.7%
Synthetic Rubber	2.3 – 2.6%	1.0%
Synthetic Fibers	5.6 – 6.3%	4.1%

Plastics and synthetic rubber materials may be further divided in the SIT into subcategories of plastics and rubber (e.g. polyethylene terephthalate (PET), polyvinyl chloride (PVC), polystyrene (PS), etc.), but the subcategories in the SIT do not match the subcategories in the waste characterization study. Therefore, the DNR did subcategorize the proportion of municipal solid waste discards.

Adjustments

As noted in the *General Calculation Method* section of this document, emissions from 2005 - 2015 were recalculated using GWPs from IPCC AR4.


Results

Total GHG emissions from the solid waste category were 1.71 MMtCO₂e in 2016, an increase of 1.00% from 2015 as shown in Table 24 and Figure 11. Emissions from municipal solid waste increased in 2016 because the cumulative amount of waste in landfills increased by 4.23% (DNR 2017).

Table 24: GHG Emissions from Municipal Solid Waste (MMtCO₂e)¹²

Pollutant	2010	2011	2012	2013	2014	2015	2016
MSW Landfills	1.45	1.45	1.66	1.48	1.46	1.68	1.70
MSW Combustion	0.02	0.02	0.02	0.02	0.02	0.01	0.01
Total	1.47	1.47	1.68	1.49	1.48	1.69	1.71

Figure 11: GHG Emissions from Solid Waste (MMtCO₂e)



¹² Totals may not equal the sum of subtotals shown in this table due to independent rounding.

Uncertainty

Excerpted from SIT Solid Waste Module (ICF 2017):

MSW Combustion

There are several sources of uncertainty in this sector, including combustion and oxidation rates, average carbon contents, and biogenic content.

- The combustion rate is not exact and varies by the quantity and composition of the waste.
- The oxidation rate varies depending on the type of waste combusted, moisture content, etc.
- The SIT uses average carbon contents instead of specific carbon contents for other plastics, synthetic rubber, and synthetic fibers.
- Non-biogenic CO₂ emissions vary depending on the amount of non-biogenic carbon in the waste and the percentage of non-biogenic carbon that is oxidized.
- The SIT assumes that all carbon in textiles is non-biomass carbon and the category of rubber and leather is almost all rubber. This may result in CO₂ emissions being slightly over-estimated (ICF 2017).

Chapter 8 – Waste: Wastewater Treatment

This chapter includes GHG emissions from the treatment of municipal and industrial wastewater. The pollutants from this sector are methane (CH₄) and nitrous oxide (N₂O). CH₄ is emitted from the treatment of wastewater, both industrial and municipal. CH₄ is produced when organic material is treated in anaerobic environment (in the absence of oxygen) and when untreated wastewater degrades anaerobically. N₂O is produced through nitrification followed by incomplete denitrification of both municipal and industrial wastewater containing both organic and inorganic nitrogen species. Production and subsequent emissions of N₂O is a complex function of biological, chemical, and physical factors, and emission rates depend on the specific conditions of the wastewater and the wastewater collection and treatment system. Human sewage makes up a significant portion of the raw material leading to N₂O emissions (ICF 2017b).

Method

Municipal Wastewater

GHG emissions from municipal wastewater are calculated in the SIT by multiplying a series of emission factors by the annual Iowa population, which was updated for 2015 (U.S. Census 2017). For example, to calculate CH₄ emissions, the state population was multiplied by the quantity of biochemical oxygen demands (BOD) per person emission factor, by the fraction that is treated anaerobically, and by the quantity of CH₄ produced per metric ton. It does not account for any digester methane that is collected and combusted instead of fossil fuels in equipment such as boilers, generators, or flares.

SIT default emission factors and assumptions were used to calculate both CH₄ and N₂O emissions, except that N₂O was calculated using the most recent protein (kg/person-year) value (45.2) from Table 7-15 in the *Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-2015* (EPA 2017b). Because the 2016 protein value was not available at the time of publication, the 2015 value was used as a surrogate for 2016.

The Iowa fraction of population without septic systems, 76%, from EPA's *Onsite Wastewater Treatment Systems Manual* (EPA 2002), was also used to estimate N₂O emissions. This value taken from the *1990 Census of Housing* and is lower than the SIT default value of 79%. The *2000 Census of Housing* and *2010 Census of Housing* do not include the Iowa fraction of population without septic systems.

Industrial Wastewater

In 2015, the DNR refined its method for calculating emissions from industrial wastewater. The DNR previously calculated emissions using the SIT and statewide red meat production numbers from the USDA. This method had a great deal of uncertainty as it only calculated emissions from wastewater at meat processing facilities and because it assumed a set amount of emissions from each metric ton of meat processed.

The EPA began requiring industrial wastewater facilities that emit 25,000 metric tons CO₂e or more to report to the federal greenhouse gas reporting program (GHGRP) starting with year 2011 emissions. In Iowa, this includes emissions from four food processing facilities and fifteen ethanol production facilities. The emissions reported to EPA have a higher level of accuracy than the SIT method because they are based on the unique characteristics and wastewater organic content of each facility. Last year nineteen ethanol production facilities and five food processing facilities emitted more than 25,000 metric tons CO₂e or more (EPA 2017a).

Adjustments

As noted in the *General Calculation Method* section of this document, emissions from 2005 - 2015 were recalculated using GWPs from IPCC AR4.

Results

Wastewater emissions account for 0.35% of total statewide GHG emissions. Total emissions from the wastewater treatment sector were 0.45 MMtCO₂e in 2016, a 0.35% increase from 2016 and a 13.70% decrease from 2005 as shown in Table 25. This is due to increases in wastewater produced by industrial meat processing facilities and the amount of municipal wastewater produced by humans as the state's population increases.


CH₄ and N₂O from municipal wastewater treatment accounted for 76.90% (0.36 MMtCO₂e) of total wastewater treatment GHG emissions as shown in Figure 12.

Table 25: GHG Emissions from Wastewater (MMtCO₂e)¹³

Sector	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Municipal CH ₄	0.236	0.237	0.239	0.240	0.241	0.244	0.245	0.246	0.248	0.249	0.250	0.251
Municipal N ₂ O	0.090	0.091	0.090	0.090	0.089	0.090	0.093	0.094	0.094	0.095	0.095	0.096
Industrial CH ₄	0.196	0.198	0.200	0.214	0.211	0.199	0.130	0.134	0.132	0.111	0.104	0.104
Total	0.522	0.526	0.529	0.543	0.541	0.533	0.468	0.473	0.474	0.455	0.449	0.451

¹³ DNR uses two decimal places throughout this report for consistency. However, in this sector three decimal places are needed show the difference in emissions from year to year.

Figure 12: GHG Emissions from Wastewater Treatment (MMtCO₂e)



*Does not include emissions from production of fruits and vegetables, pulp and paper.

Uncertainty

Excerpted from *SIT Wastewater Module (ICF 2017a)*:

Municipal Wastewater

Uncertainty is associated with both the emission factors and activity data used to calculate GHG emissions. The quantity of CH₄ emissions from wastewater treatment is based on several factors with varying degrees of uncertainty. For human sewage, there is some degree of uncertainty associated with the emission factor used to estimate the occurrence of anaerobic conditions in treatment systems based on septic tank usage data. While the Iowa-specific percentage of the population without septic systems was used to calculate emissions, the value is from 1990. There can also be variation in the per-capita BOD production association with food consumption, food waste, and disposal characteristics for organic matter. Additionally, there is variation in these factors that can be attributed to differences in wastewater treatment facilities (ICF 2017a).

N₂O emissions are dependent on nitrogen (N) inputs into the wastewater and the characteristics of wastewater treatment methods. Estimates of U.S. population, per capita protein consumption data, and the fraction of nitrogen in protein are believed to be accurate. However, the fraction that is used to represent the ratio of non-consumption nitrogen also contributes to the overall uncertainty of these calculations, as does the emission factor for effluent, which is the default emission factor from IPCC (1997). Different disposal methods of sewage sludge, such as incineration, landfilling, or land-application as fertilizer also add complexity to the GHG calculation method (ICF 2017a).

Industrial Wastewater

GHG emissions from industrial wastewater may be underestimated because only industrial wastewater facilities that emit 25,000 mtCO₂e or more are required to report to the federal greenhouse gas reporting program. Future improvements to the inventory could include identifying all of the industrial wastewater facilities that are not required to report to the federal program and developing a method to calculate their emissions.

Chapter 9 - Land Use, Land Use Change, and Forestry (LULUCF)

This chapter addresses carbon dioxide (CO₂), methane (CH₄), and nitrous oxide (N₂O) emissions from liming of agricultural soils and fertilization of settlement soils,¹⁴ as well as carbon sequestered by forests, urban trees, and yard waste and food scraps that are sent to the landfill.

Method

Forest Carbon Flux

CO₂ is taken in by plants and trees and converted to carbon in biomass during photosynthesis. “Growing forests store carbon naturally in both the wood and soil. Trees are about fifty percent carbon, and wood products from harvested trees continue to store carbon throughout their lives as well” (Flickinger 2010). CO₂ is emitted by live tree respiration, decay of dead material, fires, and biomass that is harvested and used for energy (Strait et al. 2008). The balance between the emission of carbon and the uptake of carbon is known as carbon flux (ICF 2016b).

The annual forest carbon flux was calculated using carbon storage statistics from the USDA Forest Service’s *Forest Inventory Data Online (FIDO)* (USFS 2017). FIDO data used to calculate sequestration/emission included the following forest categories:

- Carbon in live trees and saplings above ground on forest land
- Carbon in understory above ground on forest land
- Carbon in live trees and saplings below ground on forest land
- Carbon in understory below ground on forest land
- Carbon in standing dead trees on forest land
- Carbon in down dead trees on forest land
- Carbon in litter (shed vegetation decomposing above the soil surface) on forest land
- Soil organic carbon on forest land

Because 2017 carbon storage statistics were not available to calculate the 2016 carbon storage flux (2017 storage minus 2016 storage), the 2016 flux was assumed to be the same as the previous year.

Liming of Agricultural Soils

CO₂ is emitted when acidic agricultural soils are neutralized by adding limestone or dolomite. The Iowa Limestone Producers Association (ILPA) provided the DNR with the total annual amount of limestone produced for agricultural use as reported by their members (Hall 2017). However, producers do not report the percentage of limestone that is dolomitic. The Iowa Department of Transportation (DOT) tracks general information for active aggregate sources used for construction, including whether the material is limestone or dolomite. They do not track that information for limestone produced for

¹⁴ Settled soils such as landscaping, lawns, and golf courses (ICF 2017b)

agricultural purposes. The DOT indicated that some areas of the state have 100% dolomite, some have 100% limestone, and some areas are mixed (Reyes 2011). Therefore, the DNR assumed that 50% of the material produced in Iowa for agricultural use is dolomite and 50% is limestone.

Urea Fertilization

Urea emissions were calculated using the amount of urea applied annually (IDALS 2017). Because the amount of urea fertilizer applied in the in last six months of 2016 was not available, the amount applied from July 2015 – December 2015 (77,214 tons) was used as a surrogate for the amount applied from July 2016 – December 2016.

Urban Tree Flux

Carbon sequestration estimations from this sector were refined by using a new DNR data set that is a mix of land cover/remote sensing data with about a one-meter resolution. The data set includes the amount of forested acres and total acres of land for 946 incorporated areas in Iowa (Hannigan 2014).

Settlement Soils

Approximately 10% of the fertilizers applied to soils in the United States are applied to soils in settled areas such as landscaping, lawns, and golf courses (ICF 2017). N₂O emissions from settlement soils were calculated using 10% of the total annual growing year synthetic fertilizer value from the SIT Agriculture module. For more information on how the 2016 values were derived, please see *Chapter 2-Agriculture* of this report.

Non-CO₂ Emissions from Forest Fires

CH₄ and N₂O emissions from forest fires in Iowa were not estimated because the majority of wildfires and prescribed burns in Iowa that are reported to DNR occur on grasslands (Kantak 2014). In addition, the SIT calculation method uses combustion efficiencies and emission factors that are provided for primary tropical forests, secondary tropical forests, tertiary tropical forests, boreal forest, eucalypt forest, other temperate forests, shrub lands, and savanna woodlands, which are not reflective of Iowa vegetation.

Yard Trimmings and Food Scraps Stored in Landfills

GHG estimations from this sector were refined by applying the estimated percentages of yard waste and food waste in municipal solid waste from the *2011 Iowa Statewide Waste Characterization Study* (MSW 2011) to the total amount of municipal solid waste sent to landfills in 2016 (DNR 2017). While the DNR was able to use more accurate Iowa values for the annual amounts of yard waste and food scraps stored in landfills, the DNR used the SIT default values for content of yard trimmings (e.g. % grass, % leaves, % branches), carbon content, proportion of carbon stored permanently, and half-life of degradable carbon because Iowa-specific data was not available.

Adjustments

Forest Carbon Flux

The 2015 forest carbon flux was recalculated using data from the USDA Forest Service's *Forest Inventory Data Online* (USFS 2017). In the previous inventory, the 2014 carbon flux value was used as a surrogate for 2015. The previous reported value was +3.04 MMtCO₂e. The revised value is +2.87 MMtCO₂e. This changed the total emissions from the LULUCF sector in 2015 from 3.18 MMtCO₂e to 2.99 MMtCO₂e.

Urea Fertilization

Emissions from 2015 were recalculated using the most recent activity data available (IDALS 2017) as shown in Table 26. This resulted in an increase of 0.004 MMtCO₂e tons in 2015.


Table 26: Updated Urea Application (metric tons urea applied)

Year	Value Used in Dec. 2016 Report	Updated Value
2015	202,024	207,243

Results

The majority of forest carbon is stored in above ground living trees (37%) and in the forest soil (42%) as shown in Figure 13.

Figure 13: 2016 Where Forest Carbon is Stored (MMtCO₂e)




Overall, sources in the LULUCF sector released more carbon than they stored in 2016, emitting 3.13 MMtCO₂e as shown in Table 27 and Figure 14. This is an increase of 4.69% from 2015 and an increase of 115.49% from 2005. Emissions of CO₂ are shown above the x-axis in Figure 14 and carbon sinks are shown below the x-axis.

Table 27: GHG Emissions and Sinks from LULUCF (MMtCO₂e)¹⁵

Sector	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Forest Carbon Flux	-21.24	-6.53	+2.70	-4.48	-5.47	-2.68	-0.14	-0.47	-1.02	+3.04	+2.87	+2.87
Liming of Ag Soils	+0.42	+0.45	+0.37	+0.28	+0.27	+0.47	+0.51	+0.65	+0.47	+0.41	+0.34	+0.46
Urea Fertilization	+0.15	+0.15	+0.15	+0.15	+0.12	+0.11	+0.12	+0.13	+0.11	+0.15	+0.15	+0.15
Urban Trees	-0.25	-0.25	-0.25	-0.26	-0.26	-0.28	-0.28	-0.28	-0.74	-0.74	-0.74	-0.74
Yard Trimmings & Food Scraps Stored in Landfills	-0.09	-0.09	-0.08	-0.09	-0.10	-0.10	-0.13	-0.12	-0.11	-0.12	-0.12	-0.12
N ₂ O from Settlement Soils	+0.44	+0.47	+0.51	+0.48	+0.44	+0.48	+0.57	+0.57	+0.57	+0.49	+0.49	+0.51
Total	-20.56	-5.81	+3.40	-3.92	-5.00	-2.01	+0.67	+0.48	-0.71	+3.27	+2.99	+3.13

Figure 14: 2016 GHG Emissions and Sinks from LULUCF (MMtCO₂e)



¹⁵ Carbon emitted from the LULUCF sector is shown as a positive number. Carbon stored by the LULUCF sector is shown as a negative number.

Uncertainty

Uncertainty in the LULUCF sector is due to the lack of current Iowa-specific data and emission factors used to calculate emissions and/or sinks from urban trees and settlement soils. Emissions from categories such as urea fertilization, liming of agricultural soils, and yard waste and food scraps stored in landfills are more certain because Iowa-specific activity data was used. However, uncertainty was also introduced by:

- Using surrogate urea data for the last six months of 2016,
- Using growing year synthetic fertilizer data for settlement soils instead of calendar year data,
- Assuming the ratio of limestone to dolomite in Iowa is 50%, and
- Using SIT default values for content of yard trimmings (e.g. % grass, % leaves, % branches), carbon content, proportion of carbon stored permanently and half-life of degradable carbon.

In addition, due to the high uncertainty in soil carbon flux from tillage practices, it was not included in the DNR's calculations. Refer to *Chapter 2 – Agriculture* for more information.

Chapter 10 – Electricity Consumption

This chapter includes indirect emissions from electricity consumed at the point of use (e.g. residential electric hot water heaters, televisions, appliances, etc.) and does not include direct emissions generated at the electric power generating station (see Chapter 3 – Fossil Fuel Combustion).

Electricity consumed by Iowans may not be generated in Iowa. Because of this, emissions from electricity consumption do not match emissions from electricity generation (ICF 2017b). Therefore, GHG emissions from electricity consumption are included in this inventory as an informational item only and are not included in the total statewide GHG emissions to avoid any possible double counting. However, trends in electricity consumption are valuable because they are indicators of consumer behavior and trends in energy efficiency.

Method

GHG emissions were calculated using the Electricity Consumption SIT module (ICF 2017a).

Residential, Commercial, and Industrial

2016 emissions were projected by applying the forecasted percent change in energy consumption for each sector for the West North Central Region in the EIA's *Annual Energy Outlook (AEO) 2017 with Projections to 2050* (EIA 2017a) to Iowa's 2015 electricity consumption data from EIA (EIA 2017b).

Transportation

Last year for the first time DNR calculated indirect emissions from electricity consumption in the transportation sector. According to the June 2016 report *Advancing Iowa's Electric Vehicle Market* (IEDA 2016), 1,017 electric vehicles were registered in Iowa as of June 2016. This is 0.02% of the total number of vehicles, 4.41 million, registered in the state in 2016 (IDOT 2017). Emissions were calculated assuming that each electric vehicle consumes 4,250 kWh of electricity per year (IEDA 2016) and rounded to less than 0.005 MMtCO₂e as shown in Table 28. This does not include emissions from electric propulsion, other electric batteries, or non-highway electric vehicles such as golf carts.

Adjustments

2014 and 2015 emissions have been recalculated since the DNR's 2015 GHG Inventory Report was published in December 2016. The DNR previously forecasted 2015 emissions due to a lack of Iowa-specific energy consumption data. However, the 2015 energy data was released by EIA in June 2017 (EIA 2017b), so the DNR used the data to recalculate 2015 emissions as shown in Table 28 and Table 29. In addition, EPA has updated both the electricity emission factor and transmission loss factor used to calculate 2014 – 2016 emissions based on updated data in its 2014 Emissions & Generation Resource Integrated Database (eGRID) as shown in Table 28.

2005 – 2013 emissions have been updated to correct transcription errors between the calculation sheet and this TSD.

Table 28: Updated 2014 and 2015 Activity Data

Category	2014 Value Used in Dec. 2016	2014 Updated Value	2015 Value Used in Dec. 2016	2015 Updated Value
Electricity Consumption (kWh)				
Residential	14,427,000,000	14,427,251,952	13,662,369,000	13,786,000,000
Commercial	12,339,000,000	12,339,795,104	12,537,657,900	12,072,000,000
Industrial	20,436,000,000	20,437,263,558	19,765,699,200	21,289,000,000
Total	47,202,000,000	47,204,310,614	45,965,726,100	47,147,000,000
Electricity Emission Factor (lbs. CO ₂ e/kWh)	1.41	1.35	1.41	1.35
Transmission Loss Factor (%)	9.1703%	4.970%	9.1703%	4.970%

Table 29: Recalculated Electricity Emissions (MMtCO₂e)

Category	2014 Result Published in Dec. 2016	2014 Updated Result	2015 Value Published Dec. 2016	2015 Updated Result
Residential	10.16	9.29	9.62	8.88
Commercial	8.69	7.94	8.83	7.77
Industrial	14.39	13.16	13.92	13.71
Total	33.24	30.39	32.37	30.35

Results

Indirect GHG emissions from electricity consumption were 30.31 MMtCO₂e in 2016, decreasing 0.15% since 2015, mostly due to a projected 1.08% decrease in industrial electricity consumption (EIA 2017a). Industrial users consumed 44.73% of electricity in the state, while residential users consumed 29.59% and commercial users consumed 25.68% as shown in Figure 15.

Table 30: GHG Emissions from Electricity Consumption (MMtCO₂e)¹⁶

Category/Fuel Type	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Residential	12.55	12.34	12.21	12.22	10.80	11.45	11.27	9.85	10.30	9.29	8.88	8.97
Commercial	10.42	10.78	10.50	10.58	9.21	9.46	9.51	8.60	8.77	7.64	7.77	7.78
Industrial	16.56	16.95	16.61	16.71	14.33	14.84	15.14	13.74	13.83	13.16	13.71	13.56
Transportation	<i>not calculated</i>										0.00	0.00
Total	39.53	40.07	39.32	39.51	34.35	35.76	35.92	32.19	32.90	30.39	30.35	30.31

¹⁶ Totals may not equal the sum of subtotals shown in this table due to independent rounding.

Figure 15: Electricity Consumption by Category





Figure 16: Indirect GHG Emissions from Electricity Consumption by Category (MMtCO₂e)



Forecasting

Iowa Code 455B.104 requires that the DNR forecast trends in GHG emissions. The DNR projected emissions from 2016 to 2030 using the SIT Projection Tool (ICF 2017). As with many forecasts, numerous factors affect the significant level of uncertainty with future emissions. These factors may include among other things - the economy, weather, current and future environmental regulations, energy efficiency and conservation practices, driving practices, use of renewable fuels, etc.

The SIT projects that Iowa's population decreases every year from 2012 – 2030. This is contrary to the most recent population projections available from the State Data Center (Woods & Poole, 2009). Consequently, the DNR replaced the SIT default populations with the actual Iowa population for 2012 - 2015 (U.S. Census 2016) and the 2020, 2025, and 2030 projections from Woods & Poole Economics. The data points for the intervening years were calculated using a linear interpretation.

The projected emissions for 2016 – 2030 for each category are shown in Table 31. The SIT Projection Tool forecasts emissions from industrial processes, agriculture, and waste based on historical emissions from 1990 – 2014, using a combination of data sources and national projections for activity data.

Table 31: Projected Gross GHG Emissions 2016 – 2030 (MMtCO₂e)

Sector	Calculated	Projected			
	2016	2016	2020	2025	2030
Agriculture	39.48	39.01	39.70	39.54	40.02
Power Plants	25.33	27.45	30.43	26.25	23.53
RCI Fossil Fuel Use	31.90	28.39	29.97	30.61	30.77
Industrial Processes	5.50	5.47	6.44	7.71	8.89
Natural Gas Transmission & Distribution	1.41	1.18	1.21	1.30	1.30
Transportation	19.36	21.47	23.32	22.04	21.15
Waste	2.16	3.87	4.04	4.24	4.44
Total	125.15	126.82	135.11	131.69	130.11

The energy forecast is based on projected energy consumption values from the EIA's *Annual Energy Outlook (2016) with Projections to 2040* (EIA 2016a). The AEO2016 includes several different projection cases, which each address different uncertainties. The DNR used the AEO2016 "Reference Case", which represents federal and state legislation and final implementation of regulations as of the end of February 2016, including the CPP.

Short-term Projections for the Electric Power Sector

In April 2017, the U.S. Energy Information Administration of the Department of Energy announced that CO₂ emissions in the national energy sector fell 1.7% from 2015 – 2016 due to a reduction in electricity generated from coal and an increase in generation from natural gas and renewable energy. Weather also played a role, as warmer years may result in reduced energy consumption because it typically costs

less to cool a home than heat it. Heating days in the U.S. in 2016 were the second lowest of any year in the past 67 years.¹⁷

However, the most recent emissions data available for Iowa power plants does not follow the same trend as national emissions. Data from EPA's Clean Air Markets Division shows that CO₂ emissions from the electric power generation during the first six months of 2017 are 15.05% higher than CO₂ emissions from the first six months of 2016. It is unclear what caused this trend, but some factors may be a change in the number of heating and/or cooling days, changes in the prices of different fuels or other market forces.

Uncertainty

Although the SIT Projection Tool provides a good first look at projected future emissions, it has several areas of uncertainty:

1. Agricultural emissions are highly dependent on the weather and crop and livestock prices, which are not addressed by the Projection Tool.
2. In sectors where the Projection Tool predicts future emissions based on historical emissions, it only uses emissions from 1990 – 2014 and does not consider 2015 - 2016 emissions.
3. The Projection Tool forecasts emissions from fossil fuel use based on the reference case from the EIA's *Annual Energy Outlook (AEO) 2016 with Projections to 2040*, which projects emissions at the regional level and not the state level. A more current reference case, *AEO 2017*, has been published but is not used by the tool.
4. The *AEO 2016* used includes implementation of the CPP, which is currently stayed and EPA proposes to repeal.
5. The Projection Tool does not address publicly announced changes to Iowa's fossil fuel generation mix:
 - Iowa utilities have announced that from 2016 - 2025, approximately 1,000 MW of coal-fired electric generation units will retire or convert to natural gas. During that same period, approximately 185 MW of older natural gas-fired electric generation units will retire, and approximately 650 MW of newer, more efficient natural gas-fired electric generating units will come online. This will significantly reduce emissions from the electric power sector as natural gas emits approximately 50% less CO₂ per heating unit than coal emits.
 - At least 3,200 MW of additional renewables are planned for 2017 - 2020.

¹⁷ U.S. EIA: [Today in Energy - April 10, 2017](#).

References

Unless otherwise noted, all emails referenced were sent to Marnie Stein, Air Quality Bureau, Iowa Department of Natural Resources in Windsor Heights, Iowa.

General Method

ICF Consulting (2004). Emissions Inventory Improvement Program (EIIP) Volume VIII: Greenhouse Gases. Prepared for the U.S. Environmental Protection and STAPPA/ALAPCO, Washington DC.

IPCC (2001). [Climate Change 2001: Synthesis Report. A Contribution of Working Groups I, II, and III to the Third Assessment Report of the Intergovernmental Panel on Climate Change](#) [Watson, R.T. and the Core Writing Team (Eds.)]. Cambridge University Press, Cambridge, United Kingdom, and New York, New York. Available online at < >.

IPCC (2007). [Climate Change 2007: Synthesis Report. A Contribution of Working Groups I, II, and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change](#) [Core Writing Team, Pachauri, R.K. and Reisinger, A. (eds.)]. IPCC, Geneva, Switzerland.

Agriculture

Baker, J.M. et al. (2007). Tillage and soil carbon sequestration – what do we really know? *Agriculture, Ecosystems, and Environment* 118:1-5.

Blanco-Canqui, H. and R. Lal (2008). No-tillage and soil-profile carbon sequestration: an on-farm assessment. *Soil Science Society of America Journal* 72:693-701.

Boddey, R.M, C.P Jantalia, B. Alves, B. and S. Urquiaga. (2009). "Comments on 'No-Tillage and Soil-Profile Carbon Sequestration: An On-Farm Assessment'". *Soil Science Society of America Journal* 73(2):688.

DNR (2015). "Comments on EPA's Preliminary 2014 Agricultural and Grass/Pasture Burning Emissions." Correspondence from Catharine Fitzsimmons, Air Quality Bureau Chief to Venkatesh Rao and George A. Pouliot, U.S. Environmental Protection Agency, Washington DC. June 13, 2015.

EPA (2015). "[Ag Burning Emission Factors](#)". U.S. Environmental Protection Agency, Washington DC.

EPA (2016). [Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-2014](#). #430-R-16-002. U.S. Environmental Protection Agency, Washington DC.

Eve, M., D. Pape, M. Flugge, R. Steele, D. Man, M. Riley-Gilbert, and S. Biggar, Eds. (2014). [Quantifying Greenhouse Gas Fluxes in Agriculture and Forestry: Methods for Entity-Scale Inventory](#). Technical Bulletin Number 1939. Office of the Chief Economist, U.S. Department of Agriculture, Washington, DC.

Franzluebbers, A.J. (2009). "Comments on 'No-Tillage and Soil-Profile Carbon Sequestration: An On-Farm Assessment'" *Soil Science Society of America Journal* 73(2):686-7.

ICF Consulting (2004). Emissions Inventory Improvement Program (EIIP) Volume VIII: Greenhouse Gases. Prepared for the U.S. Environmental Protection and STAPPA/ALAPCO, Washington DC.

ICF International (2017a). [State Inventory Tool – Agriculture Module](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 3, 2007.

ICF International (2017b). [User’s Guide for Estimating Methane and Nitrous Oxide Emissions from Agriculture Using the State Inventory Tool](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 2017.

IDALS (2017). [Fertilizer Tonnage Distribution in Iowa for July 1, 2014 – June 30, 2015 and July 1, 2015 – June 30, 2016](#). Iowa Department of Agriculture and Land Stewardship, Commercial Feed and Fertilizer Bureau. Des Moines, Iowa.

Kantak, G. (2015). Email correspondence. Gail Kantak, Wildland Fire Supervisor, Iowa Department of Natural Resources, Des Moines, Iowa. February 6, 2015.

Licht, M. (2015). Email correspondence. Mark Licht, Cropping Systems Specialist, Iowa State University Extension, Ames, Iowa. June 6, 2015.

NRCS (2015). [Carbon Management and Evaluation Online Tool \(COMET-FARM™\)](#). National Resources and Conservation Service, Washington DC, and Colorado State University, Ft. Collins, Colorado.

Pouliot, George. (2015). Email correspondence. George Pouliot, United States Environmental Protection Agency, Washington, DC. July 30, 2015.

Stein, Marnie. (2015). Email correspondence to George Pouliot. July 30, 2015.

Strait, R. et al. (2008). [Final Iowa Greenhouse Gas Inventory and Reference Case Projections 1990 – 2025](#). Center for Climate Strategies, Washington DC.

Sucik, M. (2011a). Email correspondence. Michael Sucik, State Soil Scientist, Natural Resources and Conservation Service, Des Moines, Iowa. May 23, 2011.

Sucik, M. (2011b). Email correspondence. Michael Sucik, State Soil Scientist, Natural Resources and Conservation Service, Des Moines, Iowa. December 19, 2011.

USDA (2010). [“No-Till” Farming is a Growing Practice. Economic Information Bulletin Number 70](#). Economic Research Service, U.S. Department of Agriculture, Washington DC. Available online at < >.

USDA (2016). [2016 Iowa Agricultural Statistics Bulletin](#). National Agricultural Statistics Service, U.S. Department of Agriculture, Washington DC.

USDA (2017a). [Conservation Reserve Program – CRP Enrollment and Rental Payments by State, 1986 - 2016](#). Farm Service Agency, U.S. Department of Agriculture, Washington, DC. Accessed on August 30, 2017.

USDA (2017b). [Quick Stats 2.0: Agricultural Statistics Database](#). National Agricultural Statistics Service, U.S. Department of Agriculture, Washington DC. Accessed August 2, 2017.

West, T., and J. Six (2007). "Considering the Influence of Sequestration Duration and Carbon Saturation on Estimates of Soil Carbon Capacity". *Climatic Change*, 80(1):25-41.

Wollin, T. and W. M. Stigliani (2005). Year 2000 Iowa Greenhouse Gas Emissions Inventory. University of Northern Iowa, Cedar Falls, Iowa.

Fossil Fuel Consumption

EIA (2017a). [Annual Energy Outlook 2017 with Projections to 2050](#). Energy Information Administration, U.S. Department of Energy, Washington D.C.

EIA (2017b). [State Energy Data System \(SEDS\) 1960-2015 Completed Data File – Released June 30, 2017](#). Energy Information Administration, U.S. Department of Energy, Washington DC.

EPA (2017). [Envirofacts Greenhouse Gas Customized Search](#). U.S. Environmental Protection Agency, Washington, DC. Accessed on October 10, 2017.

ICF International (2017a). [State Inventory Tool – CO₂FFC](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 3, 2017.

ICF International (2017b). [State Inventory Tool – Stationary Combustion](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 3, 2017.

ICF International (2017c). [User's Guide for Estimating Direct Carbon Dioxide Emissions from Fossil Fuel Combustion Using the State Inventory Tool](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 2017.

ICF International (2017d). [User's Guide for Estimating Methane and Nitrous Oxide Emissions from Stationary Combustion Using the State Inventory Tool](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 2017.

IPCC/UNEP/OECD/IEA (1997). Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Intergovernmental Panel on Climate Change, United Nations Environment Programme, Organization for Economic Co-Operation and Development, International Energy Agency, Paris, France.

Industrial Processes

DNR (2010). Annual Title V Emission Inventory Data 2005 – 2009. Iowa Department of Natural Resources, Des Moines, Iowa.

EIA (2017). [Electric Power Annual 2015 – Table 2.8 Sales of Electricity to Ultimate Customers by End-Use Sector](#). Energy Information Administration, U.S. Department of Energy, Washington DC.

EPA (2016). [Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-2014](#). #430-R-16-002. U.S. Environmental Protection Agency, Washington DC.

EPA (2017a). [Envirofacts Greenhouse Gas Customized Search](#). U.S. Environmental Protection Agency, Washington, DC. Accessed on October 10, 2017.

EPA (2017b). [Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-2015. #430-P-17-001](#). U.S. Environmental Protection Agency, Washington DC.

ICF Consulting (2004). Emissions Inventory Improvement Program (EIIP) Volume VIII: Greenhouse Gases. Prepared for the U.S. Environmental Protection and STAPPA/ALAPCO, Washington DC.

ICF International (2017a). [State Inventory Tool – IP Module](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 30, 2017.

ICF International (2017b). [User’s Guide for Estimating Carbon Dioxide, Nitrous Oxide, HFC, PFC, and SF₆ Emissions from Industrial Processes Using the State Inventory Tool](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 2017.

U.S. Census Bureau (2017). [U.S. Census Quick Facts](#). U.S. Census Bureau, Washington DC. Accessed August 26, 2017.

USGS (2016). [Crushed Stone: Mineral Yearbook 2014 \[Advanced Release\]](#). Minerals Information Service, U.S. Geological Survey, Reston, Virginia.

USGS (2017). [Soda Ash: Mineral Commodity Summaries 2017](#). Minerals Information Service, U.S. Geological Survey, Reston, Virginia.

WRI (2008). [CO₂ Emissions from the Production of Ammonia v. 2.0](#). World Resources Institute Greenhouse Gas Protocol Initiative, Washington DC.

Natural Gas Transmission and Distribution

Data source: DOT (2017). [Distribution, Transmission, and Liquid Annual Data 1990 - 2016](#). Office of Pipeline Safety, Pipeline and Hazardous Materials Safety Administration, U.S. Department of Transportation. Washington DC. Accessed on August 4, 2017.

ICF International (2017a). [State Inventory Tool – Natural Gas and Oil Module](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 3, 2017.

ICF International (2017b). [User’s Guide for Estimating Carbon Dioxide and Methane Emissions from Natural Gas and Oil Systems Using the State Inventory Tool](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 2017.

Little, J. (2011). Email correspondence. Jeff Little. Energy Information Administration. June 8, 2011.

Munyon, C. (2017). Email correspondence. Cynthia Munyon, Utilities Specialist/Paralegal, Iowa Utilities Board, Des Moines, Iowa. August 7, 2017.

Transportation

EIA (2017a). [Adjusted Sales of Distillate Fuel Oil by End Use](#). U.S. Energy Information Administration. Washington DC.

EIA (2017b) [State Energy Data System \(SEDS\) 1960-2014 Completed Data File - Consumption](#). Released June 30, 2017. Energy Information Administration, U.S. Department of Energy, Washington DC.

EPA (2017). [Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-2015. #430-P-17-001](#). U.S. Environmental Protection Agency, Washington DC.

FHWA (2017). [FHWA Highway Statistics 2016 – Private and Commercial Nonhighway Use of Gasoline – 2015 \(Table MF-24\)](#). Federal Highway Administration, U.S. Department of Transportation.

ICF International (2017a). [State Inventory Tool – Mobile Combustion](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 3, 2017.

ICF International (2017b). [User’s Guide for Estimating Methane and Nitrous Oxide Emissions from Mobile Combustion Using the State Inventory Tool](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 2017.

IDOT (2017). [VMT BY County/System as of December 31 2016](#). Iowa Department of Transportation. Ames, Iowa.

Waste: Solid Waste

DNR (2017). [Tonnage Report Data](#). Iowa Department of Natural Resources, Des Moines, Iowa. Accessed on August 11, 2017.

EPA (2005). [Landfill Emission Model \(LandGEM\) Version 3.02](#). U.S. Environmental Protection Agency, Washington DC.

EPA (2011). [Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-2009](#). #430-R-11-005. U.S. Environmental Protection Agency, Washington DC.

EPA (2017). [Envirofacts Greenhouse Gas Customized Search](#). U.S. Environmental Protection Agency, Washington, DC. Accessed on October 10, 2017.

ICF International (2017). [User’s Guide for Estimating Emissions from Municipal Solid Waste Using the State Inventory Tool](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 2017.

MSW (2011). [2011 Iowa Statewide Waste Characterization Study](#). Prepared by MidAtlantic Solid Waste Consultants for the Iowa Department of Natural Resources, Des Moines, Iowa. September 2011.

Trower, B. (2017). Email correspondence. Brian Trower, Assistant Director - Electric, City of Ames, Ames, Iowa. August 8, 2017.

Waste: Wastewater Treatment

EPA (2002). [Onsite Wastewater Treatment Systems Manual. #625-R-00-008](#). U.S. Environmental Protection Agency, Washington DC.

EPA (2017a). [Envirofacts Greenhouse Gas Customized Search](#). U.S. Environmental Protection Agency, Washington, DC. Accessed on October 10, 2017.

EPA (2017b). [Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-2015. #430-P-17-001](#). U.S. Environmental Protection Agency, Washington DC.

ICF International (2017a). [State Inventory Tool – Wastewater Module](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 3, 2017.

ICF International (2017b). [User’s Guide for Estimating Methane and Nitrous Oxide Emissions from Wastewater Using the State Inventory Tool](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 2017.

U.S. Census Bureau (2017). [U.S. Census Quick Facts](#). U.S. Census Bureau, Washington DC. Accessed August 26, 2017.

LULUCF

DNR (2017). [Tonnage Report Data](#). Iowa Department of Natural Resources, Des Moines, Iowa. Accessed on August 11, 2017.

Flickinger, A. (2010). [Iowa’s Forests Today](#). Aron Flickinger, Special Projects Forester, Iowa Department of Natural Resources, Des Moines, Iowa.

Hall, J. (2017). Personal communication. Jan Hall, Iowa Limestone Producers Association, Urbandale, Iowa. August 23, 2017.

Hannigan, E. (2014). Email correspondence. Emma Hannigan, Urban Forestry Coordinator, Iowa Department of Natural Resources, Des Moines, Iowa. October 15 and 16, 2014.

ICF International (2017a). [State Inventory Tool – LULUCF Module](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 3, 2017.

ICF International (2017b). [User’s Guide for Estimating Emissions and Sinks from Land Use, Land-Use Change, and Forestry Using the State Inventory Tool](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 2017.

IDALS (2017). [Fertilizer Tonnage Distribution in Iowa for July 1, 2014 – June 30, 2015 and July 1, 2015 – June 30, 2016](#). Iowa Department of Agriculture and Land Stewardship, Commercial Feed and Fertilizer Bureau. Des Moines, Iowa.

Kantak, G. (2014). Email correspondence. Gail Kantak, Wildland Fire Supervisor, Iowa Department of Natural Resources, Des Moines, Iowa. November 10, 2014.

MSW (2011). [2011 Iowa Statewide Waste Characterization Study](#). Prepared by MidAtlantic Solid Waste Consultants for the Iowa Department of Natural Resources, Des Moines, Iowa. September 2011.

Reyes, A. (2011). Personal communication. Adriana Reyes, Geologist 3, Iowa Department of Transportation, Ames, Iowa. July 26, 2011.

Strait, R. et al. (2008). [Final Iowa Greenhouse Gas Inventory and Reference Case Projections 1990 – 2025](#). Center for Climate Strategies, Washington DC.

USFS (2017). [Forest Inventory Data Online \(FIDO\)](#). Forest Service, U.S. Department of Agriculture, Washington DC. Accessed on September 1, 2016.

Electricity Consumption

EIA (2017a). [Annual Energy Outlook 2017 with Projections to 2050](#). Energy Information Administration, U.S. Department of Energy, Washington D.C.

EIA (2017b). [State Energy Data System \(SEDS\) 1960-2015 Completed Data File – Released June 30, 2017](#). Energy Information Administration, U.S. Department of Energy, Washington DC.

ICF International (2017a). [State Inventory Tool – Electricity Consumption Module](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. April 11, 2017.

ICF International (2017b). [User’s Guide for Estimating Indirect Carbon Dioxide Equivalent Emissions from Electricity Consumption Using the State Inventory Tool](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. April 2017.

IEDA (2016). [Advancing Iowa’s Electric Vehicle Market](#). Iowa Clean Cities Coalition, Iowa Economic Development Authority, Des Moines, Iowa.

IDOT (2016). [2016 Vehicle Registrations by County](#). Iowa Department of Transportation. Ames, Iowa. Accessed on September 1, 2017.

Forecasting

CAMD (2017). [Clean Air Markets – Data and Maps](#). Clean Air Markets Division, U.S. Environmental Protection Agency, Washington D.C. Accessed on October 11, 2017.

EIA (2016a) [Annual Energy Outlook 2016 with Projections to 2040](#). Energy Information Administration, U.S. Department of Energy, Washington D.C.

EIA (2017b) [Today in Energy - April 10, 2017](#). Energy Information Administration, U.S. Department of Energy, Washington D.C.

ICF International (2017). [State Inventory Tool – Projection Tool](#). Prepared for the State Climate and Energy Program, U.S. Environmental Protection Agency, Washington DC. January 3, 2017.

U.S. Census Bureau (2017). [U.S. Census Quick Facts](#). U.S. Census Bureau, Washington DC. Accessed August 26, 2017.

Woods & Poole (2009). [Projections of Total Population for U.S., Iowa, and its Counties: 2010-2040](#). Woods & Poole Economics, Inc. 2009.

Appendix A – Iowa GHG Emissions 2005 – 2016 by Sector¹⁸

Emissions (MMtCO ₂ e)	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Agriculture	33.60	35.71	40.23	36.65	36.48	36.38	37.91	36.67	37.03	37.76	38.78	39.48
Enteric Fermentation	7.05	7.55	7.88	8.05	8.02	8.39	8.41	8.36	8.36	8.16	8.31	8.38
Manure Management	7.85	7.88	8.68	9.52	9.59	8.83	9.53	9.62	9.84	9.70	10.04	9.86
Agricultural Soil Management	18.66	20.28	23.67	19.07	18.87	19.16	19.98	18.69	18.82	19.90	20.43	21.24
Fossil Fuel Combustion	63.87	63.61	68.59	73.12	68.44	73.56	70.42	65.72	65.88	66.26	61.00	57.23
Electric Generating Facilities	36.83	36.35	40.04	41.78	37.71	42.33	38.98	35.76	33.06	33.44	29.46	25.33
Residential, Commercial, Industrial	27.03	27.26	28.55	31.34	30.73	31.23	31.44	29.96	32.82	32.82	31.54	31.90
Industrial Processes	4.57	4.71	4.78	4.92	4.22	4.79	4.49	5.17	5.34	5.28	5.28	5.50
Ammonia & Urea Production	1.01	0.91	0.95	0.87	0.60	0.84	0.75	0.85	0.88	0.86	0.81	0.92
Cement Manufacture	1.27	1.29	1.27	1.31	0.84	0.72	0.79	1.27	1.41	1.38	1.50	1.58
Electric Power Transmission & Distribution Systems	0.12	0.11	0.10	0.10	0.09	0.08	0.06	0.06	0.05	0.05	0.05	0.05
Iron and Steel Production	0.13	0.13	0.13	0.12	0.09	0.23	0.20	0.23	0.19	0.18	0.16	0.19
Lime Manufacture	0.18	0.17	0.16	0.17	0.13	0.18	0.18	0.18	0.16	0.17	0.13	0.15
Limestone and Dolomite Use	0.18	0.29	0.24	0.25	0.29	0.39	0.16	0.15	0.33	0.21	0.21	0.21
Nitric Acid Production	0.65	0.72	0.78	0.86	0.87	0.95	0.90	0.96	0.80	0.82	0.74	0.75
ODS Substitutes	1.00	1.06	1.13	1.21	1.30	1.39	1.43	1.47	1.51	1.57	1.64	1.63
Soda Ash Consumption	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
LULUCF ¹⁹	-20.56	-5.81	3.40	-3.92	-5.00	-2.01	0.67	0.48	-0.71	3.27	2.99	3.13
Forest Carbon Flux	-21.24	-6.53	2.70	-4.48	-5.47	-2.68	-0.14	-0.47	-1.02	3.04	2.87	2.87
Liming of Agricultural Soils	0.42	0.45	0.37	0.28	0.27	0.47	0.51	0.65	0.47	0.41	0.34	0.46
Urea Fertilization	0.15	0.15	0.15	0.15	0.12	0.11	0.12	0.13	0.11	0.15	0.15	0.15
Urban Trees	-0.25	-0.25	-0.25	-0.26	-0.26	-0.28	-0.28	-0.28	-0.74	-0.74	-0.74	-0.74
Yard Trimmings and Food Scraps Stored in Landfills	-0.09	-0.09	-0.08	-0.09	-0.10	-0.10	-0.13	-0.12	-0.11	-0.12	-0.12	-0.12
Fertilization of Settlement Soils	0.44	0.47	0.51	0.48	0.44	0.48	0.57	0.57	0.57	0.52	0.49	0.51

¹⁸ Totals may not equal the exact sum of subtotals in this table due to independent rounding. Values may not match values in the previous 2015 inventory published by the Department in December 2016 as they have been recalculated using the IPCC AR4 GWP. Other adjustments are described in detail in this document.

¹⁹ Carbon emitted from the LULUCF sector is shown as a positive number. Carbon stored by the LULUCF sector is shown as a negative number.

Emissions (MMtCO ₂ e)	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Natural Gas Transmission & Distribution	1.37	1.37	1.39	1.40	1.40	1.39	1.40	1.40	1.40	1.40	1.40	1.41
Transmission	0.77	0.77	0.78	0.79	0.79	0.79	0.79	0.79	0.79	0.79	0.79	0.79
Distribution	0.60	0.60	0.60	0.61	0.61	0.60	0.61	0.62	0.61	0.62	0.61	0.62
Transportation	20.14	20.35	19.96	19.99	19.40	19.41	19.58	19.59	19.46	19.55	19.31	19.36
Waste	3.10	3.03	3.07	3.10	3.06	2.01	19.94	2.15	1.96	1.94	2.14	2.16
Solid Waste	2.57	2.50	2.54	2.55	2.52	1.47	1.47	1.68	1.49	1.48	1.69	1.71
Wastewater	0.52	0.53	0.53	0.54	0.54	0.53	0.47	0.47	0.47	0.45	0.45	0.45
Gross Emissions	126.63	128.77	141.42	139.16	133.00	137.54	136.39	131.18	131.07	135.46	130.90	128.28
Sinks	-20.56	-5.81	0	-3.92	-5.00	-2.01	0	0	-0.71	0	0	0
Net Emissions	106.07	122.96	141.42	135.25	128.00	135.54	136.39	131.18	130.36	135.46	130.90	128.28
% Change from Previous Year (Gross)		+1.69%	+9.82%	-1.60%	-4.43%	+3.41%	+0.63%	-3.82%	-0.08%	+3.34	-3.37%	-2.61%
% Change from 2005 (Gross)		+1.69%	+11.68%	+9.90%	+5.03%	+8.61%	+7.71%	+3.59%	+3.51%	+6.97%	+3.37%	+1.30%

Appendix B – Iowa GHG Emissions 2005 – 2016 by Pollutant²⁰

Emissions (MMtCO ₂ e)	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Gross CO ₂	85.86	85.89	93.39	95.06	89.08	94.54	91.39	87.32	87.68	90.76	85.10	81.81
Net CO ₂	64.86	79.62	93.39	90.66	83.65	92.05	91.39	87.22	86.39	90.76	85.10	81.81
Stationary Fossil Fuel Combustion	63.54	63.28	68.25	72.76	68.09	73.09	69.96	65.30	65.47	65.85	60.64	56.94
Transportation	19.50	19.78	19.46	19.54	19.00	19.04	19.27	19.31	19.21	19.32	19.10	19.17
Industrial Processes	2.80	2.82	2.78	2.75	1.97	2.38	2.09	2.69	2.98	2.83	2.85	3.07
Solid Waste	0.03	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.01	0.01
LULUCF ²¹	-21.00	-6.28	2.89	-4.40	-5.43	-2.49	0.09	-0.09	-1.29	2.75	2.50	2.62
CH ₄	18.59	18.96	20.11	21.14	21.12	20.17	20.21	20.61	20.67	20.31	20.97	20.86
Stationary Fossil Fuel Combustion	0.10	0.10	0.10	0.10	0.11	0.19	0.19	0.17	0.17	0.17	0.14	0.10
Transportation	0.05	0.05	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.03
Natural Gas and Oil Transmission and Distribution	1.37	1.37	1.39	1.40	1.40	1.39	1.40	1.40	1.40	1.40	1.40	1.41
Enteric Fermentation	7.08	7.55	7.88	8.05	8.02	8.39	8.41	8.36	8.36	8.16	8.31	8.38
Manure Management	7.01	6.97	7.74	8.55	8.61	8.26	8.36	8.60	8.84	8.72	9.05	8.88
Solid Waste	2.55	2.49	2.52	2.54	2.50	1.45	1.45	1.66	1.48	1.46	1.68	1.70
Wastewater	0.43	0.44	0.44	0.45	0.45	0.44	0.38	0.38	0.38	0.36	0.35	0.36
N ₂ O	21.50	23.22	26.69	22.14	21.85	21.85	23.25	21.83	21.74	22.76	23.13	23.94
Stationary Fossil Fuel Combustion	0.23	0.23	0.24	0.26	0.24	0.28	0.27	0.25	0.24	0.24	0.21	0.19
Transportation	0.59	0.52	0.46	0.40	0.36	0.33	0.27	0.24	0.21	0.19	0.17	0.16
Industrial Processes	0.65	0.72	0.78	0.86	0.87	0.95	0.90	0.96	0.80	0.82	0.74	0.75
Manure Management	0.84	0.90	0.94	0.97	0.98	0.57	1.17	1.02	1.00	0.98	0.98	0.98
Agricultural Soil Management	18.66	20.28	23.67	19.07	18.87	19.16	19.98	18.69	18.82	19.90	20.43	21.24
N ₂ O from Settlement Soils	0.44	0.47	0.51	0.48	0.44	0.48	0.57	0.57	0.57	0.52	0.49	0.51
Solid Waste	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Wastewater	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.10	0.10

²⁰ Totals may not equal the exact sum of subtotals in this table due to independent rounding. Values may not match values in the previous 2015 inventory published by the Department in December 2016 as they have been recalculated using the IPCC AR4 GWP. Other adjustments are described in detail in this document.

²¹ Carbon emitted from the LULUCF sector is shown as a positive number. Carbon stored by the LULUCF sector is shown as a negative number.

Emissions (MMtCO ₂ e)	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
HFC, PFC, and SF ₆	1.12	1.17	1.23	1.31	1.39	1.47	1.49	1.53	1.56	1.62	1.69	1.68
Industrial Processes	1.12	1.17	1.23	1.31	1.39	1.47	1.49	1.53	1.56	1.62	1.69	1.68
Gross Emissions	127.07	129.24	141.42	139.65	133.44	138.02	136.39	131.28	131.65	135.46	130.90	128.28
Sinks	-21.00	-6.28	0	-4.40	-5.43	-2.49	0	-0.09	-1.29	0	0	0
Net Emissions (Sources and Sinks)	106.07	122.96	141.42	135.25	128.00	135.54	136.39	131.18	130.36	135.47	130.90	128.28